

“Cancer Reviews 2021” series

ATP and cancer immunosurveillance

Oliver Kepp^{1,2} , Lucillia Bezu^{1,2}, Takahiro Yamazaki³, Francesco Di Virgilio⁴ , Mark J Smyth⁵,
Guido Kroemer^{1,2,6,7,8,*†}  & Lorenzo Galluzzi^{3,9,10,11,12,**†} 

Abstract

While intracellular adenosine triphosphate (ATP) occupies a key position in the bioenergetic metabolism of all the cellular compartments that form the tumor microenvironment (TME), extracellular ATP operates as a potent signal transducer. The net effects of purinergic signaling on the biology of the TME depend not only on the specific receptors and cell types involved, but also on the activation status of *cis*- and *trans*-regulatory circuitries. As an additional layer of complexity, extracellular ATP is rapidly catabolized by ectonucleotidases, culminating in the accumulation of metabolites that mediate distinct biological effects. Here, we discuss the molecular and cellular mechanisms through which ATP and its degradation products influence cancer immunosurveillance, with a focus on therapeutically targetable circuitries.

Keywords ADORA2A; autophagy; CD39; CD73; immune checkpoint inhibitors; immunogenic cell death

Subject Categories Cancer; Immunology; Metabolism

DOI 10.15252/emj.2021108130 | Received 25 February 2021 | Revised 24 March 2021 | Accepted 5 April 2021 | Published online 14 June 2021

The EMBO Journal (2021) 40: e108130

Introduction

According to a widely accepted model, malignant transformation is initiated by relatively common genetic or epigenetic alterations that incapacitate tumor-suppressing mechanisms (for the most part, mechanisms that ensure the preservation of cellular homeostasis) as they activate oncogenic drivers (generally, processes that enable accrued anabolism in support of hyperproliferation) (Hanahan & Weinberg, 2011; Timp & Feinberg, 2013). The vast majority of newly

formed malignant cells, however, appears to be controlled by the host immune system prior to forming symptomatic tumors (Vesely & Schreiber, 2013; Lopez-Otin & Kroemer, 2021). While in most cases such control involves the definitive “eradication” of malignant cell precursors, in some instances newly formed cancer cells can resist immune attacks and generate a rudimentary tumor microenvironment that enables some degree of proliferation, a dynamic battle between emerging tumors and their host commonly referred to as “equilibrium” (Vesely & Schreiber, 2013). In this context, malignant cells can acquire additional genetic and epigenetic alterations that either (i) impair their ability to initiate anticancer immune responses, such as the loss or downregulation of genes coding for endogenous danger signals, (ii) limit their visibility to immune effector cells, such as the loss of MHC class I-coding genes or beta-2-microglobulin (*B2M*), (iii) increase their resistance to immune effector molecules, such as the loss of caspase 8 (*CASP8*), or (iv) establish a state of local immunosuppression, such as the upregulation of CD274 (best known as PD-L1) (Galluzzi *et al.*, 2018). In this context, the equilibrium between cancer cells and the host immune system ceases to exist in favor of an “escape” phase culminating in uncontrolled tumor growth and metastatic dissemination (Dunn *et al.*, 2002; Dersh *et al.*, 2021).

Importantly, the TME of malignancies that escaped immunosurveillance (Rao *et al.*, 2019) undergoes a considerable reconfiguration, generally involving the accumulation of immunosuppressive myeloid and lymphoid cells including M2-like tumor-associated macrophages (TAMs) and CD4⁺CD25⁺FOXP3⁺ regulatory T (T_{REG}) cells at the expense of M1-like TAMs, type I conventional dendritic cells (cDC1s), T_H1 CD4⁺ T cells, CD8⁺ cytotoxic T lymphocytes (CTLs), and natural killer (NK) cells, all of which promote tumor-targeting immunity (Talmadge & Gabrilovich, 2013; Mantovani *et al.*, 2017; Lee & Radford, 2019; Sprooten *et al.*, 2019; Togashi *et al.*, 2019). Beyond such a general trend, however, the precise immune

1 Equipe labellisée par la Ligue contre le cancer, Centre de Recherche des Cordeliers, INSERM U1138, Sorbonne Université, Université de Paris, Paris, France

2 Metabolomics and Cell Biology Platforms, Gustave Roussy Comprehensive Cancer Institute, Villejuif, France

3 Department of Radiation Oncology, Weill Cornell Medical College, New York, NY, USA

4 Department of Medical Sciences, University of Ferrara, Ferrara, Italy

5 Immunology in Cancer and Infection Laboratory, QIMR Berghofer Medical Research Institute, Herston, Brisbane, Qld, Australia

6 Pôle de Biologie, Hôpital Européen Georges Pompidou, AP-HP, Paris, France

7 Suzhou Institute for Systems Medicine, Chinese Academy of Medical Sciences, Suzhou, China

8 Department of Women’s and Children’s Health, Karolinska University Hospital, Stockholm, Sweden

9 Sandra and Edward Meyer Cancer Center, New York, NY, USA

10 Caryl and Israel Englander Institute for Precision Medicine, New York, NY, USA

11 Department of Dermatology, Yale School of Medicine, New Haven, CT, USA

12 Université de Paris, Paris, France

*Corresponding author. Tel: +33 1 4427 7667; E-mail: kroemer@orange.fr

**Corresponding author. Tel: +1 646 962 2095; E-mail: deado80@gmail.com

†These authors contributed equally to this work as senior authors

This article is part of the Cancer Reviews 2021 series.

contexture of each neoplasm exhibits considerable heterogeneity (De Sousa *et al*, 2013; Vitale *et al*, 2021) and has a major impact on disease course and response to therapy (Fridman *et al*, 2017). Indeed, it has now become clear that the efficacy of most anticancer agents commonly employed in the clinic, encompassing cytotoxic chemotherapeutics, radiation therapy (RT), and targeted anticancer agents, relies at least partially on the (re)activation of immunosurveillance (Galluzzi *et al*, 2020; Rodriguez-Ruiz *et al*, 2020; Petroni *et al*, 2021). In line with this notion, considerable efforts are being dedicated to the identification of clinically relevant approaches to alter the TME in favor of treatment efficacy, especially for tumors that exhibit rather scarce infiltration by immune effector cells, such as luminal breast cancer and pancreatic carcinoma (Kroemer *et al*, 2015; Ho *et al*, 2020).

All cellular components of the TME including malignant and immune cells engage in a dynamic competition for nutrients, oxygen, and trophic signals (all of which are generally scarce as a consequence of relatively poor vascularization) (Martinez-Outschoorn *et al*, 2017; O'Sullivan *et al*, 2019; Garner & de Visser, 2020). Moreover, the availability of nutrients, oxygen, and trophic signals is not equal across all tumor regions and is not constant over time (e.g., before and after therapy), hence constituting a major driver of intratumoral heterogeneity (ITH) (De Sousa *et al*, 2013; Vitale *et al*, 2021). Indeed, such restrictions *de facto* operate as Darwinian pressures, fostering the selection of cells with an accrued capacity to harness alternative carbon sources (e.g., lactate, extracellular amino acids) for catabolic and anabolic reactions in support of proliferation and tolerance to hypoxia (Chang *et al*, 2015; Xiao *et al*, 2019).


Adenosine triphosphate (ATP) occupies a key position in the overall configuration of the TME. On one hand, intracellular ATP is crucial for each cellular TME component to survive and mediate its functions (including proliferation, for malignant and non-terminally differentiated immune cells) (Leone & Powell, 2020; Bergers & Fendt, 2021). On the other hand, the pool of ATP that accesses the TME upon active secretion by living or dying cells into the extracellular space constitutes a major signal transducer (Di Virgilio *et al*, 2018). The net effect of ATP signaling on the TME, however, depends on multiple factors, including the presence of extracellular ATP-degrading enzymes as well as the expression pattern of purinergic receptors. Here, we will critically discuss the molecular and cellular mechanisms through which extracellular ATP and its degradation products influence the crosstalk between malignant and immune cells and present recent advances on the purinergic system as a potential target for the development of novel anticancer interventions.

Extracellular ATP homeostasis in the TME

Since ATP cannot be synthesized in the extracellular milieu, the microenvironmental levels of ATP are entirely controlled by the balance between its secretion/release and degradation (Fig 1).

Adenosine triphosphate secretion is an active, regulated process that can occur via multiple mechanisms and involve different cellular sources. Molecular mechanisms for ATP secretion encompass: (i) exocytosis of ATP-containing vesicles, a process that may (but does not necessarily) involve cell death (Imura *et al*, 2013; Martins

et al, 2014) and mechanistically relies on vesicular loading by solute carrier family 17 member 9 (SLC17A9) (Imura *et al*, 2013; Cao *et al*, 2014) and the SNAP receptor (SNARE)- and Rho-associated, coiled-coil containing protein kinase 1 (ROCK1)-dependent fusion of exocytosis-competent ATP-rich vesicles with the plasma membrane (Imura *et al*, 2013; Martins *et al*, 2014); (ii) liberation of cytosolic ATP molecules via gap junction protein alpha 1 (GJA1, best known as CX43) hemichannels at gap junctions (Stout *et al*, 2002; Eltzschig *et al*, 2006; Kang *et al*, 2008); and (iii) gradient-driven efflux via oligomeric pannexin 1 (PANX1) channels (also known as “pannexons”) (Dahl, 2015), which are sensitive to activation by mechanical forces (Bao *et al*, 2004), by the pro-inflammatory CASP1 (Narahari *et al*, 2021), and by apoptotic caspases such as CASP3 and CASP7 (Chekeni *et al*, 2010; Medina *et al*, 2020). That said, while both vesicular ATP secretion and PANX1-dependent release have been documented in living and dying malignant cells (Martins *et al*, 2014; Martin *et al*, 2017), CX43 hemichannels appear to be mostly operational in non-malignant cells of the TME, including astrocytes (Stout *et al*, 2002), as well as (at least potentially) neutrophils (Dosch *et al*, 2018) and macrophages (Dosch *et al*, 2019). Indeed, while both neutrophils and macrophages have been shown to release ATP via CX43 hemichannels in non-oncological settings, whether such function is preserved in TAMs and tumor-infiltrating neutrophils (TINs) remains to be elucidated. Along similar lines, platelets are known as major sources of extracellular ATP upon degranulation (Yeaman, 2014), but their contribution to extracellular ATP availability in the TME has just begun to emerge (Schumacher *et al*, 2013; Gaertner & Massberg, 2019). Additional cellular compartments that may secrete ATP in the TME encompass (at least in some settings) endothelial cells (Sáez *et al*, 2018; Yang *et al*, 2020d), fibroblasts (Pinheiro *et al*, 2013; Murata *et al*, 2014), dendritic cells (DCs) (Tappe *et al*, 2018; Martinek *et al*, 2019), and activated CTLs (Tokunaga *et al*, 2010). Importantly, while some cells spontaneously secrete at least some ATP in their physiological status, for the most part, ATP is actively released in the context of adaptive responses to microenvironmental perturbations, including mechanical cues (Bao *et al*, 2004), inflammatory signals (Beckel *et al*, 2018), hypoxia (Lim To *et al*, 2015), and exposure to a variety of cancer therapeutics (Michaud *et al*, 2011; Tatsuno *et al*, 2019; Rodriguez-Ruiz *et al*, 2020). In most such instances, abundant ATP secretion by stressed cells (which is key for extracellular ATP to mediate immunostimulatory effects, see below) involves functional autophagic responses (Michaud *et al*, 2011), potentially linked to the ability of autophagy to preserve intracellular ATP pools during stress (Rybstein *et al*, 2018; Anderson & Macleod, 2019). Consistent with this notion, genetic and pharmacological interventions aimed at blocking or boosting autophagic responses in cancer cells have been consistently associated with reduced and increased ATP secretion, respectively, in response to immunogenic chemotherapy (Michaud *et al*, 2011; Pietrocola *et al*, 2016; Chen *et al*, 2019; Kepp & Kroemer, 2020; Wang *et al*, 2020). Obviously, all dying cells abruptly release their cytosolic ATP pool when they undergo plasma membrane permeabilization (PMP) as the final step of cellular demise. However, while PMP itself has now been shown to be an active (rather than an osmosis-driven) process even in the context of post-apoptotic, secondary necrosis (Kayagaki *et al*, 2021), the consequent spillage of cytosolic content into the extracellular milieu remains a largely unregulated phenomenon.



© EMBO

Figure 1. Extracellular ATP homeostasis in the tumor microenvironment.

The concentration of extracellular ATP in the tumor microenvironment is determined by the balance between ATP release and degradation. A variety of cells release ATP either as part of their physiological state or as they respond to stress and potentially die, including cancer cells, dendritic cells (DCs), tumor-infiltrating neutrophils (TINs), tumor-associated macrophages (TAMs), and platelets. Extracellular ATP is catabolized by the sequential activity of ectonucleoside triphosphate diphosphohydrolase 1 (ENTPD1, best known as CD39), which converts ATP into ADP and AMP, and 5'-nucleotidase ecto (NTSE, best known as CD73), which converts AMP into adenosine (ADO). CD39 and CD73 are expressed by multiple cell type that populate the tumor microenvironment, including some malignant cells, cancer-associated fibroblasts (CAFs), exhausted cytotoxic T lymphocytes (CTLs), regulatory T (T_{REG}) cells, an immunosuppressive subset of natural killer (NK) cells, M2-like TAMs, and myeloid-derived suppressor cells (MDSCs). Hypoxia is a major driver of ATP degradation in the tumor microenvironment.

Extracellular ATP is rapidly catalyzed by the sequential activity of two ectonucleotidases, that is, ectonucleoside triphosphate diphosphohydrolase 1 (ENTPD1, best known as CD39), which converts ATP into ADP and AMP, and 5'-nucleotidase ecto (NT5E, best known as CD73), which converts AMP into adenosine as the rate-limiting step of this enzymatic cascade (Allard *et al*, 2020; Moesta *et al*, 2020). CD39 is mostly expressed by T_{REG} cells (Borsellino *et al*, 2007), M2-like TAMs (d'Almeida *et al*, 2016), and myeloid-derived suppressor cells (MDSCs, an immature population of myeloid cells with potent immunosuppressive activity) (Li *et al*, 2017), as well as by specific cancer cell types, such as adult T-cell leukemia/lymphoma cells (Nagate *et al*, 2021), type 16 human papillomavirus (HPV-16)-associated cervical carcinoma cells (de Lourdes Mora-García *et al*, 2019), and ovarian carcinoma cells (Häusler *et al*, 2011). Moreover, CD8⁺ CTLs undergoing terminal exhaustion as a consequence of chronic antigen stimulation generally exhibit a CD39⁺ phenotype (Canale *et al*, 2018). Conversely, CD73 is expressed by a wide variety of malignant cells as well as by cancer-associated fibroblasts (CAFs) (Yu *et al*, 2020), T_{REG} cells (Stagg *et al*, 2011), and a regulatory subset of NK cells (Neo *et al*, 2020). Interestingly, the whole-body deletion of purinergic receptor P2X 7 (P2RX7), which codes for one of the main receptors of extracellular ATP, has a major impact on extracellular ATP levels in the TME of experimental P2RX7-competent melanomas (De Marchi *et al*, 2019), at least in part as a consequence of altered tumor infiltration by CD39⁺ and CD73⁺ T_{REG} cells and decreased ATP release by TAMs (De Marchi *et al*, 2019). Such an effect, however, cannot be recapitulated by the pharmacological P2RX7 antagonist A740003 as a result of its mixed activity on immune cells (it fails to alter tumor infiltration by T_{REG} cells, decreases the abundance of intratumoral CD39⁺ and CD73⁺ effector T (T_{EFF}) cells, and inhibits ATP secretion by TAMs) and malignant cells (it favors ATP secretion by malignant cells) (De Marchi *et al*, 2019).

Of note, extracellular ATP degradation does not necessarily require the expression of CD39 and CD73 on the same cell (*in cis*), but can also occur efficiently when these ectonucleotidases are expressed by different cellular compartments that are in proximity to each other within the TME (Schuler *et al*, 2014). CD73 is abundant in T_{REG} cell-derived exosomes (Smyth *et al*, 2013), which are highly mobile and hence further promote the overall catalytic efficiency of ATP degradation within the TME. Moreover, CD38 (also known as cyclic ADP-ribose hydrolase) and ectonucleotide pyrophosphatase/phosphodiesterase 1 (ENPP1), which are expressed by some cancer cells and exhausted T cells, can compensate for limited CD39 activity as they catalyze the conversion of extracellular NAD⁺ into ADP ribose and AMP (Morandi *et al*, 2015). Finally, the expression of both CD39 and CD73 can be upregulated by hypoxia, which is relatively common in the TME of solid neoplasms, via a transcriptional mechanism that involves hypoxia-inducible factor 1 subunit alpha (HIF1A, best known as HIF-1 α) (Giatromanolaki *et al*, 2020; Synnestvedt *et al*, 2002).

In summary, the levels of extracellular ATP in the TME are dynamically determined by the mutually opposed inputs of secretion/release *vs.* degradation. As the factors governing these aspects of the ATP biology exhibit a considerable degree of ITH, regional and temporal fluctuations in extracellular ATP levels are likely to play a major role in the outcome of purinergic signaling in the TME, as discussed further below.

Immunostimulation by extracellular ATP

Extracellular ATP mediates two main functions: (i) It operates as a chemotactic cue for myeloid cells, upon binding to the purinergic receptor P2Y2 (P2RY2), a metabotropic receptor (Elliott *et al*, 2009; Chekeni *et al*, 2010), and (ii) it promotes activation of the inflammasome and hence CASP1-dependent secretion of interleukin 1 beta (IL1B) and IL18 upon binding to P2RX7, an ionotropic receptor (Péregaux *et al*, 2000). Importantly, both these effects are required for the optimal activation of tumor-specific immune responses by (and hence the complete efficacy of) immunogenic chemotherapeutics such as anthracyclines and oxaliplatin, as demonstrated in *P2ry2*^{-/-}, *P2rx7*^{-/-}, *Casp1*^{-/-} and *Il18*^{-/-} mice, as well as mice lacking a core component of the inflammasome (*Nlrp3*^{-/-} mice), the main IL1B receptor (*Il1r1*^{-/-} mice) or treated with a purinergic receptor antagonist (suramin) or an IL1B-blocking antibody (Ghiringhelli *et al*, 2009; Aymeric *et al*, 2010; Ma *et al*, 2013).

In this setting, DC precursors newly recruited to the TME via ATP released by cancer cells succumbing to immunogenic cell death (ICD) not only mature upon ATP-driven inflammasome activation and migrate to tumor-draining lymph nodes or tertiary lymphoid structures to prime adaptive anticancer immunity, but also recruit a population of IL17-producing $\gamma\delta$ T cells that is critical for tumor infiltration by primed CTLs (Ma *et al*, 2011). In accordance with this notion, optimal anticancer immune responses (and consequent superior therapeutic efficacy) driven by immunogenic chemotherapeutics are compromised in *Il17a*^{-/-} and *Il17ra*^{-/-} mice (Ma *et al*, 2011). Intriguingly, it has recently shown that the chemotactic activity of ATP on DCs also involves P2RX7 and PANX1 (Saez *et al*, 2017), suggesting the existence of a feed-forward loop whereby intracellular ATP stores may contribute to DC migratory capacity (Saez *et al*, 2017). Moreover, elevated levels of extracellular ATP appear to induce pyroptosis in P2RX7⁺ M2-like TAMs, hence supporting T cell-mediated antitumor immunity upon the depletion of immunosuppressive cells from the TME (Bidula *et al*, 2019).

Importantly, the net immunomodulatory effect of extracellular ATP depends on the activation of additional signaling pathways. Indeed, the PANX1-dependent co-release of ATP and a wide panel of metabolites including ADP, AMP, GMP, creatine, spermidine, and glycerol-3-phosphate (G3P) by dying cells reportedly promote the removal of cell corpses while preventing the initiation of inflammatory reactions (Medina *et al*, 2020; Narahari *et al*, 2021). Moreover, extracellular ATP can have direct tumorigenic functions. Specifically, the cancer cell-driven release of ATP from platelets initiates a P2RY2-dependent signaling cascade that promotes tumor extravasation and metastatic dissemination upon the opening of endothelial barriers (Schumacher *et al*, 2013; Chen *et al*, 2019; Wang *et al*, 2020). The autophagy-dependent secretion of ATP by melanoma cells has been shown to promote invasiveness and resistance to the BRAF inhibitor vemurafenib, a process that requires P2RX7 expression in the malignant cell compartment (Martin *et al*, 2017). Similar findings have been obtained with human triple-negative breast cancer (TNBC) MDA-MB-231 cells upon the ATP-dependent activation of the transcription factor SRY-box transcription factor 9 (SOX9) (Yang *et al*, 2020a). Finally, NME/NM23 nucleoside diphosphate kinase 1 (NME1, best known as NDPK-A) and NME2 (best known as NDPK-B) expression on extracellular vesicles from MDA-

MB-231 cells reportedly support the formation of pulmonary metastatic niches as a consequence of extracellular ATP generation *in situ* and consequent activation of purinergic receptor P2Y1 (P2RY1) (Duan *et al*, 2021).

Consistent with the multipronged effects of extracellular ATP on the TME, a large body of clinical literature suggests that genetic or epigenetic defects affecting ATP signaling influence disease outcome

in cancer patients in a context-dependent manner (Table 1). For instance, while loss-of-function polymorphisms in *P2RX7* (rs3751143; rs208294) have been associated with advanced stage or poor disease outcome in cohorts of patients with breast carcinoma (Ghiringhelli *et al*, 2009), chronic lymphocytic leukemia (CLL) (Thunberg *et al*, 2002; Wiley *et al*, 2002; Zhang *et al*, 2003), and papillary thyroid carcinoma (PTC) (Dardano *et al*, 2009), no impact on

Table 1. Pathophysiological relevance of extracellular ATP signaling in human cancer.

Cancer	No. patients	Variable	Technology	Impact	References
Breast cancer	225	<i>P2RX7</i> rs3751143	SNP analysis	Metastatic dissemination Decreased OS	Ghiringhelli <i>et al</i> (2009)
	1,067 1,992	<i>BECN1</i>	Gene expression profiling	Improved disease outcome	Tang <i>et al</i> (2015)
	152	<i>MAP1LC3B</i>	IHC	Improved MFS Improved OS	Ladoire <i>et al</i> (2016)
	152 1,646	<i>MAP1LC3B</i>	IHC	Improved PFS	Ladoire <i>et al</i> (2015)
CLL	36	<i>P2RX7</i> rs3751143	SNP analysis	Disease stage	Wiley <i>et al</i> (2002)
	144	<i>P2RX7</i> rs3751143	SNP analysis	Marginally decreased OS	Zhang <i>et al</i> (2003)
	170	<i>P2RX7</i> rs3751143	SNP analysis	Decreased OS	Thunberg <i>et al</i> (2002)
	111	<i>P2RX7</i> rs3751143	SNP analysis	No correlation with clinical outcome	Nüchel <i>et al</i> (2004)
	121	<i>P2RX7</i> rs3751143	SNP analysis	No correlation with clinical outcome	Starczynski <i>et al</i> (2003)
	21	<i>P2RX7</i>	Immunoblotting	Disease progression	Adinolfi <i>et al</i> (2002)
Colorectal cancer	2,297	<i>MAP1LC3B</i>	Gene expression profiling	Increased OS	Li <i>et al</i> (2020)
Gastric cancer	14	<i>P2RY2</i>	Gene expression profiling	Disease	Aquea <i>et al</i> (2014)
	354	<i>MAP1LC3C</i>	Gene expression profiling	Improved OS	Wang <i>et al</i> (2021)
	354	<i>ATG4D</i>	Gene expression profiling	Decreased OS	Wang <i>et al</i> (2021)
	402	<i>MAP1LC3B</i> and <i>SQSTM1</i>	IHC, immunoblotting and RT-PCR	Decreased OS	Kim <i>et al</i> (2019)
Head and neck cancer	79	<i>MAP1LC3B</i>	IHC	Disease stage	Jiang <i>et al</i> (2012)
Hepatocellular carcinoma	190	<i>MAP1LC3B</i>	IHC	Improved OS	Lee <i>et al</i> (2013)
	1,086	<i>BECN1</i>	Gene expression profiling	Improved OS	Qin <i>et al</i> (2018)
Multiple myeloma	136	<i>P2RX7</i> rs3751143	SNP analysis	No correlation with clinical outcome	Paneesha <i>et al</i> (2006)
Ovarian cancer	1,497	<i>BECN1</i>	Gene expression profiling	Improved OS and PFS	Chen <i>et al</i> (2020c)
	1,497	<i>MAP1LC3B</i>	Gene expression profiling	No correlation with clinical outcome	Chen <i>et al</i> (2020c)
Pancreatic cancer	73	<i>BECN1</i>	IHC	Disease progression	Ko <i>et al</i> (2013)
	86	<i>BECN1</i> and <i>MAP1LC3B</i>	IHC and RT-PCR	Metastatic dissemination Disease stage Decreased OS	Cui <i>et al</i> (2019)
Papillary thyroid cancer	121	<i>P2RX7</i> rs3751143	SNP analysis	Disease type	Dardano <i>et al</i> (2009)
	121	<i>P2RX7</i> rs208294	SNP analysis	No effect on disease stage	Dardano <i>et al</i> (2009)
Salivary gland carcinoma	48	<i>BECN1</i> and <i>MAP1LC3B</i>	IHC	Low disease stage	Li <i>et al</i> (2019a)

CLL, chronic lymphocytic leukemia; IHC, immunohistochemistry; MFS, metastasis-free survival; N/A, not available; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism.

clinicopathological variables could be attributed to rs3751143 in other cohorts of subjects with CLL (Starczyński *et al*, 2003; Nüchel *et al*, 2004), multiple myeloma (Paneesha *et al*, 2006), and PTC (Dardano *et al*, 2009), while increased expression levels of P2RX7 have been linked to disease progression in an independent cohort of CLL patients (Adinolfi *et al*, 2002). Likewise, elevated P2RY2 levels have been associated with gastric malignant transformation (Aquea *et al*, 2014). A variety of immunohistochemical and transcriptional signatures of proficient autophagic responses have been linked to worsened disease outcome in cohorts of breast (Yamazaki *et al*, 2020), gastric (Kim *et al*, 2019; Wang *et al*, 2021), pancreatic (Ko *et al*, 2013; Cui *et al*, 2019), and head and neck (Jiang *et al*, 2012) cancer patients, while the contrary held true (or there was no impact on clinicopathological variables) in independent series of patients with breast (Ladoire *et al*, 2015; Tang *et al*, 2015; Ladoire *et al*, 2016), ovarian (Chen *et al*, 2020c), hepatocellular (Lee *et al*, 2013; Qin *et al*, 2018), gastric (Wang *et al*, 2021), colorectal (Li *et al*, 2020), and salivary gland (Li *et al*, 2019a) carcinoma.

In summary, while an abundant preclinical literature mechanistically implicates ATP secretion by stressed and dying cells in the initiation of anticancer immune responses (Fig 2), additional, hitherto poorly characterized factors and mechanisms appear to influence the net effect of ATP signaling in the TME.

Immunosuppression by extracellular ATP metabolites

Extracellular ATP degradation by the sequential activity of CD39 and CD73 mediates immunosuppressive effects not only as a consequence of limited ATP-dependent immunostimulation (see above), but also due to the generation of adenosine, which *per se* promotes tumor progression via immunological and non-immunological mechanisms (Allard *et al*, 2020). Consistent with this notion, transgene-driven overexpression of CD39 by malignant cells has been associated with impaired anticancer immunity driven by immunogenic chemotherapy (and hence poor disease outcome) in syngeneic immunocompetent mouse models of fibrosarcoma (Michaud *et al*, 2011; Pietrocola *et al*, 2016). Similarly, experimental CD73 overexpression in human cervical carcinoma as well as human and mouse breast carcinoma cells has been shown to promote invasiveness and metastatic potential (Zhou *et al*, 2007; Gao *et al*, 2017), at least in part as a consequence of autocrine–paracrine adenosine signaling via adenosine A2b receptor (ADORA2B) and ADORA2B-driven neovascularization (Stagg *et al*, 2010; Mittal *et al*, 2016; Ludwig *et al*, 2020). Together with ADORA2A, ADORA2B is indeed the main receptor for extracellular adenosine signaling in the TME (Allard *et al*, 2020). However, while ADORA2B is expressed by various malignant cell types including breast cancer (Stagg *et al*, 2010; Lan *et al*, 2018), cervical cancer (Torres-Pineda *et al*, 2020), and melanoma (Mittal *et al*, 2016) cells, as well as by endothelial cells (Ludwig *et al*, 2020), DCs (Chen *et al*, 2020b), and M2-like TAMs (Cohen *et al*, 2015), ADORA2A expression is mostly restricted to myeloid cells (Nakamura *et al*, 2020), NK cells (Young *et al*, 2018), CTLs (Kjaergaard *et al*, 2018; Shi *et al*, 2019), and lymphatic endothelial cell precursors (Allard *et al*, 2019).

From a mechanistic standpoint, ADORA2A- and ADORA2B-driven immunosuppression results from the activation of intracellular cyclic AMP (cAMP) signaling, the ultimate functional outcome of

which depends on the specific immune cell type expressing these receptors. Thus, cAMP signaling directly inhibits TCR activation and cytokine production in CTLs (Lappas *et al*, 2005), promotes forkhead box P3 (FOXP3) synthesis and the upregulation of the co-inhibitory receptor cytotoxic T lymphocyte-associated protein 4 (CTLA4) in T_{REG} cells (Klein & Bopp, 2016), impairs NK cell effector and secretory functions (Raskovalova *et al*, 2005), inhibits NF- κ B-dependent inflammatory responses (Minguet *et al*, 2005) and drives the secretion of immunosuppressive cytokines such as IL10 from TAMs and MDSCs (Németh *et al*, 2005; Cekic *et al*, 2014), and perturbs the maturation of DCs (Kayhan *et al*, 2019). Of note, both ADORA2A and ADORA2B can be upregulated by hypoxia (Sitkovsky *et al*, 2014; Lan *et al*, 2018), in thus far resembling CD39 and CD73. Thus, particularly hypoxic regions of the TME are expected to exhibit robust adenosinergic signaling and hence (i) potent immunosuppression, (ii) accrued neovascularization, (iii) increased vascular permeability, and (iv) enhanced cancer cell motility, *de facto* representing ideal niches for metastatic dissemination. This is further aggravated by immunometabolic circuitries driven by hypoxia. Specifically, hypoxic tumor regions are rich in glycolytic lactate, which mediates multipronged immunosuppressive effects involving CTLs (Brand *et al*, 2016), NK cells (Husain *et al*, 2013), T_{REG} cells (Watson *et al*, 2021), and MDSCs (Yang *et al*, 2020c).

Consistent with the potent immunosuppressive effects of adenosine signaling in the TME, an abundant clinical literature links elevated expression levels of adenosine-generating enzymes or adenosine receptors to poor disease progression in cohorts of patients with various tumors (Table 2). Thus, high levels of CD39 have been associated with advanced grade or poor disease outcome in multiple cohorts of patients with CLL (Pulte *et al*, 2011), renal cell carcinoma (Wu *et al*, 2020a), endometrial tumors (Aliagas *et al*, 2014), pancreatic carcinoma (Künzli *et al*, 2007), and non-small cell lung cancer (NSCLC; Li *et al*, 2017). Similar clinical findings have been correlated with various single nucleotide polymorphisms affecting *Entpd1* (i.e., rs10748643, rs11188513, rs2226163) in patients with colorectal carcinoma (Tokunaga *et al*, 2019; Gallerano *et al*, 2020), but the functional impact of these variants on CD39 functions remains unclear. Moreover, abundant tumor-infiltrating or circulating levels CD4⁺ or CD8⁺ cells expressing CD39 have been linked to disease progression or resistance to therapy in various cohorts of individuals with CLL (Perry *et al*, 2012), colorectal carcinoma (Gallerano *et al*, 2020), head and neck squamous cell carcinoma (Gallerano *et al*, 2020), pancreatic cancer (Gallerano *et al*, 2020), NSCLC (Koh *et al*, 2020), and renal cell carcinoma (Qi *et al*, 2020). Along these lines, intratumoral or circulating biomarkers of CD73 proficiency (including expression levels and enzymatic activity) have been correlated with poor disease outcome in patients with diffuse large B-cell lymphoma (Wang *et al*, 2019), ovarian cancer (Turcotte *et al*, 2015), pancreatic carcinoma (Chen *et al*, 2020a; Tahkola *et al*, 2021), NSCLC (Li *et al*, 2017), renal cell carcinoma (Tripathi *et al*, 2020), breast cancer (Loi *et al*, 2013; Buisseret *et al*, 2018), glioma (Xu *et al*, 2013), colorectal carcinoma (Messaudi *et al*, 2020), and metastatic melanoma (Turiello *et al*, 2020). The *Nt5e* polymorphism rs2229523 (of hitherto unclear functional significance) correlated with limited overall survival in patients with colorectal carcinoma (Tokunaga *et al*, 2019), as did tumor infiltration by CD8⁺CD73⁺ cells in prostate cancer patients (Leclerc *et al*, 2016). Finally, ADORA2A expression or signaling was linked to poor




Figure 2. Immunostimulation by extracellular ATP.

Extracellular ATP mediates chemotactic effects on myeloid cells, upon binding to purinergic receptor P2Y2 (P2RY2), and promotes the inflammasome-dependent secretion of interleukin 1 beta (IL1B) and IL18 upon binding to purinergic receptor P2X7 (P2RX7). These effects are critical for mature dendritic cells (DCs) to prime cytotoxic T lymphocytes (CTLs) against tumor-derived antigens and hence initiate adaptive anticancer immunity. Alongside, extracellular ATP triggers pyroptosis in tumor-associated macrophages (TAMs), hence depleting the tumor microenvironment of generally immunosuppressive cells along with the emission of pyroptosis-dependent immunostimulatory signals.

disease outcome in individuals with diffuse large B-cell lymphoma (Wang *et al*, 2019), renal carcinoma (Kamai *et al*, 2021), and a variety of tumors from The Cancer Genome Atlas (Sidders *et al*, 2020), as was the *ADORA2B* polymorphism rs2015353 in patients with colorectal cancer (Tokunaga *et al*, 2019), although the functional implications of rs2015353 on *ADORA2B* activity remain to be elucidated.

Of note, the discovery that the TME contains unusually high levels of extracellular ATP has prompted innovative therapeutic approaches that specifically harness this biochemical feature. For instance, a monoclonal antibody specific for the immunostimulatory receptor TNF receptor superfamily member 9 (TNFRSF9, best known as CD137) has been engineered to drive CD137 signaling only in the presence of nearly millimolar ATP levels, thus inducing a potent anticancer immune response in the absence of adverse effects due to extratumoral activity (Kamata-Sakurai *et al*, 2020). This agent is currently being tested in combination with the immune checkpoint inhibitor (ICI) atezolizumab (a PD-L1-blocking antibody) for safety and preliminary efficacy in a phase I clinical trial enrolling patients with solid tumors (JapicCTI-205153). Along similar lines, it has recently been shown that a novel positive allosteric modulator of P2RX7 (which only acts in the presence of high extracellular ATP levels) potentiates the therapeutic effects of an ICI specific for programmed cell death 1 (PDCD1, best known as PD-1) by stimulating DCs to release IL18 in support of the effector functions of tumor-

infiltrating NK cells and CTLs (Douguet *et al*, 2021). Incidentally, these therapeutic applications provide an independent and convincing demonstration (*mutatis mutandis*, almost an “*ex adjuvantibus*” proof) of the accuracy of early measurements of extracellular ATP concentration in the TME (Pellegatti *et al*, 2008).

Taken together, these observations point to CD39/CD73-dependent adenosine generation and consequent adenosinergic signaling via *ADORA2A* and *ADORA2B* as a key immunosuppressive mechanism supporting the progression and metastatic dissemination multiple tumors (Fig 3). Of note, while additional adenosine receptors including *ADORA1* and *ADORA3* (which inhibit cAMP signaling) are expressed by malignant cells and some tumor-infiltrating immune cells (Stagg & Smyth, 2010), their role in immunosurveillance remains poorly investigated.

Targeting purinergic signaling for cancer therapy

Consistent with the key role of extracellular ATP and its degradation products in the control of immunosurveillance, a variety of pharmacological and genetic approaches designed to boost ATP-driven immunostimulation or inhibit adenosine-dependent immunosuppression (alone or combined with other treatments) have been shown to mediate prominent antineoplastic effects in immunocompetent mouse models of cancer.

Table 2. Pathophysiological relevance of adenosinergic signaling in human cancer.

Cancer	No. patients	Variable	Technology	Impact	References
Breast cancer	122	CD73	IHC	Decreased OS	Buisseret et al (2018)
	>6,000	<i>NT5E</i>	Gene expression profiling	Reduced pCR	Loi et al (2013)
CLL	34	Circulating CD4 ⁺ CD39 ⁺ and CD8 ⁺ CD39 ⁺ cells	Flow cytometry	Disease stage	Pulte et al (2011)
	62	Circulating CD4 ⁺ CD39 ⁺ T cells	Flow cytometry	Disease stage	Perry et al (2012)
Colorectal cancer	107	<i>ADORA2B</i> rs2015353	SNP analysis	Decreased OS	Tokunaga et al (2019)
	107 215	<i>ENTPD1</i> rs11188513	SNP analysis	Decreased OS	Tokunaga et al (2019)
	215	<i>ENTPD1</i> rs2226163	SNP analysis	Decreased OS	Tokunaga et al (2019)
	107	<i>NT5E</i> rs2229523	SNP analysis	Decreased OS	Tokunaga et al (2019)
	129	<i>ENTPD1</i> , <i>NT5E</i> , and <i>ADORA2B</i>	SNP analysis	No correlation	Tokunaga et al (2019)
	60	Circulating CD8 ⁺ CD39 ⁺ T cells	Flow cytometry	Disease progression	Gallerano et al (2020)
	60	<i>ENTPD1</i> rs10748643	SNP analysis	Disease progression	Gallerano et al (2020)
	215	CD73	IHC	Disease progression Decreased OS	Messaoudi et al (2020)
	193	Circulating CD73	ELISA	Decreased OS	Messaoudi et al (2020)
DLBCL	91	<i>ADORA2A</i> on TILs	IHC	Decreased OS	Wang et al (2019)
	91	CD73	IHC	Decreased OS	Wang et al (2019)
Endometrial tumors	29	CD39	IHC	Tumor grade	Aliagas et al (2014)
Glioma	500	<i>NT5E</i>	Gene expression profiling	Limited DFS	Xu et al (2013)
HNSCC	19	Circulating CD8 ⁺ CD39 ⁺ T cells	Flow cytometry	Disease progression	Gallerano et al (2020)
Melanoma	546	Circulating CD73	AMPase activity	Disease progression Decreased OS	Turiello et al (2020)
NSCLC	132	Circulating CD8 ⁺ CD39 ⁺ T cells	Flow cytometry	Decreased OS	Koh et al (2020)
	24	Circulating CD8 ⁺ CD39 ⁺ MDSCs	Flow cytometry	Tumor infiltration by MDSCs	Li et al (2017)
Ovarian cancer	208	CD73	IHC	Decreased OS	Turcotte et al (2015)
Pancreatic cancer	28	<i>ENTPD1</i>	RT-PCR	Decreased OS	Künzli et al (2007)
	3	Circulating CD8 ⁺ CD39 ⁺ T cells	Flow cytometry	Disease progression	Gallerano et al (2020)
	110	CD73	IHC	Decreased OS	Tahkola et al (2021)
	168	<i>NT5E</i>	Gene expression profiling	Disease progression Decreased OS	Chen et al (2020a)
Prostate cancer	285	Circulating CD8 ⁺ CD73 ⁺ T cells	IHC	Disease progression	Leclerc et al (2016)
Renal cell carcinoma	60	<i>ADORA2A</i> expression	IHC	Metastatic dissemination Decreased OS	Kamai et al (2021)
	138	CD73	IHC	Decreased OS	Tripathi et al (2020)
	243	CD8 ⁺ CD39 ⁺ T cells	IHC	Disease progression	Qi et al (2020)
	367	CD39	IHC and RT-PCR	Disease stage Disease progression	Wu et al (2020a)
Various tumor types	N/A	<i>ADORA2A</i> -regulated gene expression	Gene expression profiling	Decreased OS	Sidders et al (2020)

CLL, chronic lymphocytic leukemia; DLBCL, diffuse large B-cell lymphoma; DFS, disease-free survival; ELISA, enzyme-linked immunosorbent assay; IHC, immunohistochemistry; MDSC, myeloid-derived suppressor cell; N/A, not available; OS, overall survival; pCR, pathological complete response; SNP, single nucleotide polymorphism; TIL, tumor-infiltrating lymphocyte.




Figure 3. Immunosuppression by extracellular ATP metabolites.

Extracellular adenosine (ADO) not only favors metastatic dissemination by binding adenosine A2b receptor (ADORA2B) on malignant cells, hence promoting invasiveness, and endothelial cells (ECs), thus promoting neovascularization, but also modulates the functions of multiple immune cells upon interaction with ADORA2B or adenosine A2a receptor (ADORA2A). Specifically, adenosine signaling inhibits dendritic cell (DC) maturation and interferes with the effector functions of cytotoxic T lymphocytes (CTLs), B cells, and natural killer (NK) cells, while favoring immunosuppression by M2-like tumor-associated macrophages (TAMs), myeloid-derived suppressor cells (MDSCs), and regulatory T (T_{REG}) cells.

Boosting ATP secretion with autophagy-activating maneuvers, including weekly cycles of nutrient deprivation and administration of so-called caloric restriction mimetics (CRMs, i.e., molecules that induce autophagy and cause other biochemical correlates of nutrient deprivation in the absence of sizeable weight loss), has been linked to improved therapeutic responses to immunogenic chemotherapies in immunocompetent mouse models of fibrosarcoma, correlating with decreased infiltration by T_{REG} cells, via a mechanism that depends on expression of the essential autophagy gene autophagy-related 5 (*ATG5*) in cancer cells and intact immune responses (Pietrocola *et al*, 2016; Castoldi *et al*, 2020; Wu *et al*, 2020b). Along similar lines, short-term starvation reportedly boosts the responsiveness of mouse breast cancer cells growing in immunocompetent syngeneic hosts to radiation therapy (RT) (Saleh *et al*, 2013; Simone *et al*, 2016). However, proficient autophagic responses in mouse breast cancer cells limit the efficacy of RT *in vivo* as a consequence of an improved disposal of permeabilized mitochondria that would otherwise release mitochondrial DNA in the cytosol and trigger cyclic GMP-AMP synthase (CGAS) signaling coupled to type I interferon (IFN) secretion (Medler *et al*, 2019; Yamazaki *et al*, 2020). This suggests that whole-body autophagy activation by short-term fasting may support the efficacy of RT by mechanisms unrelated to ATP secretion in the TME, potentially linked to improved autophagic responses in immune cells, most of which rely on autophagy for optimal functions (Clarke & Simon, 2019). That said, even though cyclic/short-term nutrient deprivation could be safely implemented in at least some cancer patients (Krstic *et al*, 2020), clinicians remain cautious on implementing clinical trials involving such a nutritional measure. Similarly, while many CRMs with robust

anticancer activity in preclinical models are currently available as over-the-counter medications (e.g., aspirin) (Castoldi *et al*, 2020) or vitamin supplementations (e.g., nicotinamide) (Buqué *et al*, 2020), clinical development remains at bay, at least in some cases reflecting potential dosing issues.

Blocking CD39 or CD73 has also been associated with robust antineoplastic effects in immunocompetent mouse models of cancer. For instance, whole-body *Entpd1* deletion as well as reconstitution of radiosensitive hematopoietic cells with *Entpd1*^{-/-} precursors have been shown to inhibit *in vivo* growth and metastatic dissemination of mouse melanoma and colorectal carcinoma cells (Sun *et al*, 2010; Perrot *et al*, 2019). Similar findings have been obtained in *Entpd1*^{-/-} mice xenografted with mouse melanoma or fibrosarcoma cells and treated with immunogenic chemotherapy or ICIs (Perrot *et al*, 2019), as well as with (i) monoclonal antibodies targeting human CD39 in human *ENTPD1* knock-in mice bearing mouse fibrosarcoma cells (Perrot *et al*, 2019), (ii) monoclonal antibodies targeting mouse CD39 in wild-type mice used as hosts for mouse melanoma, fibrosarcoma, or colorectal carcinoma cells (Li *et al*, 2019c; Yan *et al*, 2020), (iii) antisense oligonucleotides targeting *Entpd1* in wild-type mice bearing syngeneic breast cancer cells (Kashyap *et al*, 2019), and (iv) pharmacological CD39 inhibitors in wild-type mice xenografted with mouse colorectal carcinoma cells (Michaud *et al*, 2011). Of note, in the latter models, the anticancer effects of CD39 blockage could be abrogated by *P2rx7* or *Nlrp3* whole-body deletion, as well as by co-administration of monoclonal antibodies targeting IL18 (Li *et al*, 2019c; Yan *et al*, 2020), formally linking therapeutic activity to accrued extracellular ATP signaling (rather than to mere adenosine depletion). Mice lacking *Nt5e* have

Table 3. Ongoing clinical trials targeting purinergic signaling for cancer therapy.^a

Agent	Target	Indications	Phase	Status	Notes	Ref.
AZD4635 (AstraZeneca)	ADORA2A	CRPC	II	Recruiting	In combination with oleclumab (anti-CD73), durvalumab (anti-PD-L1)	NCT04089553
		NSCLC	I/II	Active	In combination with oleclumab (anti-CD73)	NCT03381274
		Solid tumors	I	Active	Single agent and in combination with durvalumab (anti-PD-L1), oleclumab (anti-CD73), docetaxel, abiraterone acetate, enzalutamide	NCT02740985
		Solid tumors	I	Recruiting	Single agent	NCT03980821
Ciforadenant CPI-444/V81444 (Corvus Pharmaceuticals)	ADORA2A	NSCLC	I/II	Recruiting	Single agent and in combination with atezolizumab (anti-PD-L1)	NCT03337698
		CPRC Renal cell carcinoma	I	Recruiting	Single agent and in combination with atezolizumab (anti-PD-L1)	NCT02655822
EOS100850 (iTeos Therapeutics)	ADORA2A	Solid tumors	I	Recruiting	Single agent	NCT03873883
Etrumadenant AB928 (Arcus Biosciences)	ADORA2A ADORA2B	CRC GEC	I	Active	In combination with mFOLFOX	NCT03720678
		NSCLC	I/I	Recruiting	In combination with carboplatin, pemetrexed and pembrolizumab	NCT03846310
	Ovarian cancer TNBC	I	Recruiting	In combination with IPI-549 (PI3γ inhibitor), doxorubicin, paclitaxel	NCT03719326	
	Solid tumors	I	Active	In combination with zimberelimab (AB122, anti-PD1)	NCT03629756	
Taminadenant NIR178/PBF-509 (Pablobio/Novartis)	ADORA2A	NSCLC	I	Active	Single agent and in combination with spartalizumab (anti-PD-1)	NCT02403193
		TNBC	I	Recruiting	In combination with spartalizumab (anti-PD-1) and LAG525 (anti-LAG-3)	NCT03742349
		Solid tumors	I	Recruiting	Single agent and in combination with NZV930 (anti-CD73), spartalizumab (anti-PD1)	NCT03549000
		Solid tumors	I	Recruiting	Single agent and in combination with spartalizumab (anti-PD-1); NZV930 (anti-CD73), KAZ954	NCT04237649
		Solid tumors	II	Recruiting	Single agent and in combination with spartalizumab (anti-PD-1)	NCT03207867
SRF617 (Surface Oncology)	CD39	Solid tumors	I	Recruiting	Single agent and in combination with paclitaxel, gemcitabine, pembrolizumab (anti-PD-L1)	NCT04336098
TTX-030 (AbbVie)	CD39	Lymphoma Solid tumors	I/I	Recruiting	Pembrolizumab (anti-PD-L1), docetaxel, gemcitabine	NCT03884556
		Solid tumors	I	Recruiting	In combination with budigalimab (anti-PD-1), mFOLFOX	NCT04306900
BMS986179 (Bristol Myers Squibb)	CD73	Solid tumors	I/II	Active	Single agent and in combination with nivolumab (anti-PD-1)	NCT02754141
CPI-006 (Corvus Pharmaceuticals)	CD73	NHL Solid tumors	I/I	Recruiting	Single agent and in combination with ciforadenant (ADORA2A antagonist), pembrolizumab (anti-PD-L1)	NCT03454451
LY3475070 (Eli-Lilly)	CD73	Solid tumors	I	Recruiting	Single agent or in combination with pembrolizumab (anti-PD-L1)	NCT04148937
NZV930 (Surface Oncology)	CD73	Solid tumors	I	Recruiting	Single agent or in combination with spartalizumab (anti-PD-1), NIR178 (ADORA2A antagonist)	NCT03549000
Oleclumab MEDI9447 (AstraZeneca)	CD73	Bladder cancer	I	Recruiting	In combination with durvalumab (anti-PD-L1)	NCT03773666
		NSCLC	II	Recruiting	In combination with durvalumab (anti-PD-L1)	NCT03334617
		Ovarian cancer	II	Recruiting	Single agent or in combination with durvalumab (anti-PD-L1)	NCT03267589
		Pancreatic cancer	I/II	Recruiting	In combination with gemcitabine, paclitaxel, durvalumab (anti-PD-L1), FOLFOX	NCT03611556
		TNBC	I/II	Recruiting	In combination with durvalumab (anti-PD-L1), paclitaxel	NCT03742102

Table 3 (continued)

Agent	Target	Indications	Phase	Status	Notes	Ref.
TJ004309 TJD5 (Tracon Pharmaceuticals)	CD73	Solid tumors	I	Recruiting	In combination with atezolizumab (anti-PD-L1)	NCT03835949

CRC, colorectal cancer; CRPC, castration-resistant prostate cancer; GEC, gastroesophageal cancer; NHL, non-Hodgkin lymphoma; NSCLC, non-small cell lung cancer; TNBC, triple-negative breast cancer.

^aRestricted to active and recruiting studies, as per www.clinicaltrials.gov on February 15, 2021.

been shown to be poorly permissive to the growth of syngeneic glioblastoma, lymphoma, melanoma, ovarian cancer, colorectal carcinoma, and breast cancer cells (Stagg *et al*, 2011; Wang *et al*, 2011; Yegutkin *et al*, 2011; Young *et al*, 2016; Yan *et al*, 2019). Moreover, blockage of CD73 with monoclonal antibodies or pharmacological agents mediates standalone therapeutic effects or enhances the efficacy of other treatments (encompassing chemotherapeutics, RT, and ICIs) in immunocompetent mouse models of breast cancer (Loi *et al*, 2013; Allard *et al*, 2014; Young *et al*, 2016; Wennerberg *et al*, 2020), colorectal carcinoma (Allard *et al*, 2013; Hay *et al*, 2016; Tsukui *et al*, 2020), melanoma (Iannone *et al*, 2014; Young *et al*, 2016), ovarian cancer (Häusler *et al*, 2014; Li *et al*, 2019b), head and neck squamous cell carcinoma (Deng *et al*, 2018), and prostate cancer (Allard *et al*, 2013). In line with such an abundant preclinical literature, several monoclonal antibodies targeting CD39 or CD73 are currently being tested for their safety and anticancer efficacy, either as standalone therapeutics or combined with ICIs, in clinical trials (Table 3). Preliminary findings from such studies point to an acceptable safety profile and promising clinical activity encompassing disease stabilization and (at least in some patients) partial or complete responses (Mobasher *et al*, 2019; Bendell *et al*, 2020).

Genetic and pharmacological strategies for ADORA2A and ADORA2B inhibition have also been shown to mediate anticancer effects in preclinical cancer models. For instance, (whole-body or myeloid cell-specific) *Adora2a* or *Adora2b* deletion has been demonstrated to inhibit tumor growth in immunocompetent mice bearing syngeneic melanoma (Ohta *et al*, 2006; Cekic *et al*, 2014; Chen *et al*, 2020b), lymphoma (Waickman *et al*, 2012; Nakamura *et al*, 2020), breast carcinoma (Beavis *et al*, 2013), or lung cancer cells (Chen *et al*, 2020b). Consistent with this notion, pharmacological ADORA2A (e.g., CPI-444, SCH58261, AZD4635) or ADORA2B (e.g., PSB-1115) inhibitors have been attributed robust antineoplastic properties in syngeneic mouse models of melanoma (Willingham *et al*, 2018), fibrosarcoma (Beavis *et al*, 2017), breast cancer (Beavis *et al*, 2013; Mittal *et al*, 2014; Beavis *et al*, 2015; Beavis *et al*, 2017), lymphoma (Nakamura *et al*, 2020), colorectal carcinoma (Beavis *et al*, 2015; Willingham *et al*, 2018), multiple myeloma (Yang *et al*, 2020b), and renal cell carcinoma (Willingham *et al*, 2018), especially when combined with ICIs or other immunotherapies. Importantly, at least in some of these models, co-inhibition of ADORA2A and CD73 mediated superior tumor control as compared to inhibiting ADORA2A or CD73 alone (Young *et al*, 2016), suggesting a non-redundant role for these two factors in the establishment of local immunosuppression. Consistent with these preclinical observations, a number of ADORA2A or dual ADORA2A/ADORA2B inhibitors are currently being investigated for safety and activity in clinical trials (Table 3). Preliminary findings from these studies indicate that

many ADORA2A/ADORA2B inhibitors exhibit an acceptable safety profile and at least some degree of clinical activity (Chiappori *et al*, 2018; Fong *et al*, 2020; Lim *et al*, 2020).

Additional approaches that may be harnessed to target adenosine signaling in the TME encompass the use of CD38 or ENPP1 blockers and strategies that revert tumor hypoxia (e.g., respiratory hyperoxygenation) (Hatfield *et al*, 2014; Hatfield & Sitkovsky, 2020). However, CD38-specific agents (including the FDA-approved monoclonal antibody daratumumab) are currently used with the aim of eradicating CD38-expressing myeloma cells (Facon *et al*, 2019), and ENPP1 blockers are still in preclinical development (Carozza *et al*, 2020). Similarly, hyperoxygenation has been employed in the past to improve radiosensitivity (because the ability of radiation therapy to cause DNA damage in cancer cell depends on local oxygen tension) but is no longer used for this purpose.

In summary, although a variety of strategies have been successfully used to target purinergic signaling in preclinical tumor models, CD73, ADORA2A, and (less so) CD39 blockers are the only drugs currently in clinical development as anticancer agents, for the most part in combination with standard of care therapeutics or ICIs (Table 3).

Concluding remarks

In summary, extracellular ATP and its degradation products play a key role in the regulation of the tumor immune contexture, hence have a major influence on the propensity of human neoplasms to respond to therapy. While various agents aimed at boosting extracellular ATP concentrations and/or limiting adenosine signaling are already in clinical development, multiple questions to be addressed and avenues to be explored remain. First, it will be important to determine the contribution of P2RX7 and/or P2RY2 signaling in cancer cells to tumor growth and metastatic dissemination in specific settings. Accumulating evidence indicates that some cancer cells can harness extracellular ATP in support of disease progression and resistance to therapy (Martin *et al*, 2017), suggesting that CD39 and/or CD73 inhibition may not mediate optimal anticancer effects in some settings, and calling for the identification of cancer cell-targeted P2RX7 or P2RY2 inhibitors. Second, it will be crucial to develop combinatorial approaches based on the dual blockage of ATP degradation and adenosine signaling, potentially in the context of ICI-based immunotherapy. Indeed, it appears that the efficacy of ADORA2A and/or ADORA2B blockers can be significantly boosted by CD73 inhibition (Young *et al*, 2016), pointing to a functional non-redundancy that may be harnessed for therapeutic purposes. Also, while inhibiting ATP degradation or adenosine signaling mediates anticancer effects *per se* (at least in various models), blocking

additional immunosuppressive pathways such as those controlled by co-inhibitory receptors appears to provide improved disease control in most cases (Beavis *et al*, 2017; Leone *et al*, 2018; Goswami *et al*, 2020). Finally, it will be interesting to elucidate whether and how purinergic signaling can be targeted in patients with innate or acquired resistance to ICI-based immunotherapy, reflecting the fact that exhausted T cells generally express high levels of CD39 (Canale *et al*, 2018). Irrespective of these and other incognita, purinergic signaling stands out as a particularly promising target for the development of novel anticancer agents.

Acknowledgements

We are indebted to Giulia Petroni (Weill Cornell Medicine) for help with figure preparation. FDV is supported by the Italian Association for Cancer Research (AIRC) grant # IG13025 and by institutional funds from the University of Ferrara (Italy). MJS is supported by a National Health and Medical Research Council Investigator (1173958) and Program Grant (1132519). GK is supported by the Ligue contre le Cancer (équipe labellisée); Agence National de la Recherche (ANR)—Projets blancs; ANR under the frame of E-Rare-2, the ERA-Net for Research on Rare Diseases; Association pour la recherche sur le cancer (ARC); Association Ruban Rose; Cancéropôle Ile-de-France; Fondation pour la Recherche Médicale (FRM); a donation by Elior; Gustave Roussy Odyssey, the European Union Horizon 2020 Project Oncobiome; Fondation Carrefour; High-end Foreign Expert Program in China (GDW20171100085 and GDW20181100051), Institut National du Cancer (INCa); Inserm (HTE); Institut Universitaire de France; LeDucq Foundation; the LabEx Immuno-Oncology; the RHU Torino Lumière; the Seerave Foundation; the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); and the SIRIC Cancer Research and Personalized Medicine (CARPEM). The LG laboratory is supported by a Breakthrough Level 2 grant from the US Department of Defense (DoD), Breast Cancer Research Program (BRCP) (#BC180476P1), by the 2019 Laura Ziskin Prize in Translational Research (#ZP-6177, PI: Formenti) from the Stand Up to Cancer (SU2C), by a Mantle Cell Lymphoma Research Initiative (MCL-RI, PI: Chen-Kiang) grant from the Leukemia and Lymphoma Society (LLS), by a startup grant from the Dept. of Radiation Oncology at Weill Cornell Medicine (New York, US), by a Rapid Response Grant from the Functional Genomics Initiative (New York, US), by industrial collaborations with Lytix (Oslo, Norway) and Phosplatin (New York, US), and by donations from Phosplatin (New York, US), the Luke Heller TECPR2 Foundation (Boston, US), and Sotio a.s. (Prague, Czech Republic).

Author contributions

LG conceived the article. OK and LG wrote the first version of the manuscript with critical input from LB, TY, FDV, MJS, and GK. OK, TY, and LB prepared display items under supervision from LG. All authors approved the final version of the article.

Conflict of interest

OK reports research funding from Daiichi Sankyo and a co-founder role of Samsara therapeutics. FDV is a member of the Scientific Advisory Board of Biosceptre, Ltd. GK reports research funding from Bayer Healthcare, Eleor, Genentech, Glaxo Smyth Kline, Institut Mérieux, Lytix, PharmaMar, Sotio, and Vasculox (completed), funding from Samsara, consulting/advisory honoraria from The Longevity Labs and Lytix, membership of the Executive Board of Bristol Myers Squibb Foundation France, co-founder role of everImmune, Samsara therapeutics and Therfast-Bio. LG reports research funding from Lytix and Phosplatin (completed), consulting/advisory honoraria from

Boehringer Ingelheim, AstraZeneca, OmniSEQ, The Longevity Labs, Inzen, Onxeo, and the Luke Heller TECPR2 Foundation. All other authors declare that they have no conflict of interest.

References

- Adinolfi E, Melchiorri L, Falzoni S, Chiozzi P, Morelli A, Tieghi A, Cuneo A, Castoldi G, Di Virgilio F, Baricordi OR (2002) P2X7 receptor expression in evolutive and indolent forms of chronic B lymphocytic leukemia. *Blood* 99: 706–708
- Aliagas E, Vidal A, Texidó L, Ponce J, Condom E, Martín-Satué M (2014) High expression of ecto-nucleotidases CD39 and CD73 in human endometrial tumors. *Mediators Inflamm* 2014: 509027
- Allard B, Allard D, Buisseret L, Stagg J (2020) The adenosine pathway in immuno-oncology. *Nat Rev Clin Oncol* 17: 611–629
- Allard B, Cousineau I, Allard D, Buisseret L, Pommey S, Chrobak P, Stagg J (2019) Adenosine A2a receptor promotes lymphangiogenesis and lymph node metastasis. *Oncimmunology* 8: 1601481
- Allard B, Pommey S, Smyth MJ, Stagg J (2013) Targeting CD73 enhances the antitumor activity of anti-PD-1 and anti-CTLA-4 mAbs. *Clin Cancer Res* 19: 5626–5635
- Allard B, Turcotte M, Spring K, Pommey S, Royal I, Stagg J (2014) Anti-CD73 therapy impairs tumor angiogenesis. *Int J Cancer* 134: 1466–1473
- Anderson CM, Macleod KF (2019) Autophagy and cancer cell metabolism. *Int Rev Cell Mol Biol* 347: 145–190
- Aquea G, Bresky G, Lancellotti D, Madariaga JA, Zaffiri V, Urzua U, Haberle S, Bernal G (2014) Increased expression of P2RY2, CD248 and EphB1 in gastric cancers from Chilean patients. *Asian Pac J Cancer Prev* 15: 1931–1936
- Aymeric L, Apetoh L, Ghiringhelli F, Tesniere A, Martins I, Kroemer G, Smyth MJ, Zitvogel L (2010) Tumor cell death and ATP release prime dendritic cells and efficient anticancer immunity. *Cancer Res* 70: 855–858
- Bao L, Locovei S, Dahl G (2004) Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett* 572: 65–68
- Beavis PA, Divisekera U, Paget C, Chow MT, John LB, Devaud C, Dwyer K, Stagg J, Smyth MJ, Darcy PK (2013) Blockade of A2A receptors potently suppresses the metastasis of CD73+ tumors. *Proc Natl Acad Sci USA* 110: 14711–14716
- Beavis PA, Henderson MA, Giuffrida L, Mills JK, Sek K, Cross RS, Davenport AJ, John LB, Mardiana S, Slaney CY *et al* (2017) Targeting the adenosine 2A receptor enhances chimeric antigen receptor T cell efficacy. *J Clin Invest* 127: 929–941
- Beavis PA, Milenkovski N, Henderson MA, John LB, Allard B, Loi S, Kershaw MH, Stagg J, Darcy PK (2015) Adenosine receptor 2A blockade increases the efficacy of anti-PD-1 through enhanced antitumor T-cell responses. *Cancer Immunol Res* 3: 506–517
- Beckel JM, Gómez NM, Lu W, Campagno KE, Nabet B, Albalawi F, Lim JC, Boesze-Battaglia K, Mitchell CH (2018) Stimulation of TLR3 triggers release of lysosomal ATP in astrocytes and epithelial cells that requires TRPML1 channels. *Sci Rep* 8: 5726
- Bendell JC, Manji GA, Pant S, Lai DW, Colabella J, Berry W, Paoloni MC, Grossman WJ, O'Reilly EM (2020) A phase I study to evaluate the safety and tolerability of AB680 combination therapy in participants with gastrointestinal malignancies. *J Clin Oncol* 38: TPS788
- Bergers G, Fendt SM (2021) The metabolism of cancer cells during metastasis. *Nat Rev Cancer* 21: 162–180

- Bidula S, Dhuna K, Helliwell R, Stokes L (2019) Positive allosteric modulation of P2X7 promotes apoptotic cell death over lytic cell death responses in macrophages. *Cell Death Dis* 10: 882
- Borsellino G, Kleinewietfeld M, Di Mitri D, Sternjak A, Diamantini A, Giometto R, Höpner S, Centonze D, Bernardi G, Dell'Acqua ML et al (2007) Expression of ectonucleotidase CD39 by Foxp3⁺ Treg cells: hydrolysis of extracellular ATP and immune suppression. *Blood* 110: 1225–1232
- Brand A, Singer K, Koehl G, Kolitzus M, Schoenhammer G, Thiel A, Matos C, Bruss C, Klobuch S, Peter K et al (2016) LDHA-associated lactic acid production blunts tumor immunosurveillance by T and NK cells. *Cell Metab* 24: 657–671
- Buisseret L, Pommey S, Allard B, Garaud S, Bergeron M, Cousineau I, Ameyle L, Bareche Y, Paesmans M, Crown J et al (2018) Clinical significance of CD73 in triple-negative breast cancer: multiplex analysis of a phase III clinical trial. *Ann Oncol* 29: 1056–1062
- Buqué A, Bloy N, Perez-Lanzón M, Iribarren K, Humeau J, Pol JG, Levesque S, Mondragon L, Yamazaki T, Sato Ai et al (2020) Immunoprophylactic and immunotherapeutic control of hormone receptor-positive breast cancer. *Nat Commun* 11: 3819
- Canale FP, Ramello MC, Núñez N, Araujo Furlan CL, Bossio SN, Gorosito Serrán M, Tosello Boari J, Del Castillo A, Ledesma M, Sedlik C et al (2018) CD39 expression defines cell exhaustion in tumor-infiltrating CD8(+) T cells. *Cancer Res* 78: 115–128
- Cao Q, Zhao K, Zhong XZ, Zou Y, Yu H, Huang P, Xu TL, Dong XP (2014) SLC17A9 protein functions as a lysosomal ATP transporter and regulates cell viability. *J Biol Chem* 289: 23189–23199
- Carozza JA, Brown JA, Bohnert V, Fernandez D, AlSaif Y, Mardjuki RE, Smith M, Li L (2020) Structure-aided development of small-molecule inhibitors of ENPP1, the extracellular phosphodiesterase of the immunotransmitter cGAMP. *Cell Chem Biol* 27: 1347–1358
- Castoldi F, Humeau J, Martins I, Lachkar S, Loew D, Dingli F, Durand S, Enot D, Bossut N, Chery A et al (2020) Autophagy-mediated metabolic effects of aspirin. *Cell Death Discov* 6: 129
- Cekic C, Day YJ, Sag D, Linden J (2014) Myeloid expression of adenosine A2A receptor suppresses T and NK cell responses in the solid tumor microenvironment. *Cancer Res* 74: 7250–7259
- Chang C-H, Qiu J, O'Sullivan D, Buck M, Noguchi T, Curtis J, Chen Q, Gindin M, Gubin M, van der Windt G et al (2015) Metabolic competition in the tumor microenvironment is a driver of cancer progression. *Cell* 162: 1229–1241
- Chekeni FB, Elliott MR, Sandilos JK, Walk SF, Kinchen JM, Lazarowski ER, Armstrong AJ, Penuela S, Laird DW, Salvesen GS et al (2010) Pannexin 1 channels mediate 'find-me' signal release and membrane permeability during apoptosis. *Nature* 467: 863–867
- Chen G, Xie W, Nah J, Sauvat A, Liu P, Pietrocola F, Sica V, Carmona-Gutierrez D, Zimmermann A, Pendl T et al (2019) 3,4-Dimethoxychalcone induces autophagy through activation of the transcription factors TFE3 and TFEB. *EMBO Mol Med* 11: e10469
- Chen Q, Pu N, Yin H, Zhang J, Zhao G, Lou W, Wu W (2020a) CD73 acts as a prognostic biomarker and promotes progression and immune escape in pancreatic cancer. *J Cell Mol Med* 24: 8674–8686
- Chen S, Akdemir I, Fan J, Linden J, Zhang B, Cekic C (2020b) The expression of adenosine A2B receptor on antigen-presenting cells suppresses CD8(+) T-cell responses and promotes tumor growth. *Cancer Immunol Res* 8: 1064–1074
- Chen X, Sun Y, Wang B, Wang H (2020c) Prognostic significance of autophagy-related genes Beclin1 and LC3 in ovarian cancer: a meta-analysis. *J Int Med Res* 48: 300060520968299
- Chiappori A, Williams CC, Creelan BC, Tanvetyanon T, Gray JE, Haura EB, Thapa R, Chen D-T, Beg AA, Boyle TA et al (2018) Phase I/II study of the A2AR antagonist NIR178 (PBF-509), an oral immunotherapy, in patients (pts) with advanced NSCLC. *J Clin Oncol* 36: 9089
- Clarke AJ, Simon AK (2019) Autophagy in the renewal, differentiation and homeostasis of immune cells. *Nat Rev Immunol* 19: 170–183
- Cohen HB, Ward A, Hamidzadeh K, Ravid K, Mosser DM (2015) IFN- γ prevents adenosine receptor (A2bR) upregulation to sustain the macrophage activation response. *J Immunol* 195: 3828–3837
- Cui L, Wang X, Zhao X, Kong C, Li Z, Liu Y, Jiang X, Zhang X (2019) The autophagy-related genes Beclin1 and LC3 in the prognosis of pancreatic cancer. *Int J Clin Exp Pathol* 12: 2989–2996
- d'Almeida SM, Kauffenstein G, Roy C, Basset L, Papargyris L, Henrion D, Catros V, Ifrah N, Descamps P, Croue A et al (2016) The ecto-ATPase CD39 is involved in the acquisition of the immunoregulatory phenotype by M-CSF-macrophages and ovarian cancer tumor-associated macrophages: regulatory role of IL-27. *Oncoimmunology* 5: e1178025
- Dahl G (2015) ATP release through pannexon channels. *Philos Trans R Soc Lond B Biol Sci* 370: 20140191
- Dardano A, Falzoni S, Caraccio N, Polini A, Tognini S, Solini A, Berti P, Di Virgilio F, Monzani F (2009) 1513A>C polymorphism in the P2X7 receptor gene in patients with papillary thyroid cancer: correlation with histological variants and clinical parameters. *J Clin Endocrinol Metab* 94: 695–698
- de Lourdes Mora-García M, López-Cisneros S, Gutiérrez-Serrano V, García-Rocha R, Weiss-Steider B, Hernández-Montes J, Sánchez-Peña HI, Ávila-Ibarra LR, Don-López CA, Muñoz-Godínez R et al (2019) HPV-16 Infection is associated with a high content of CD39 and CD73 Ectonucleotidases in cervical samples from patients with CIN-1. *Mediators Inflamm* 2019: 4651627
- De Marchi E, Orioli E, Pegoraro A, Sangaletti S, Portararo P, Curti A, Colombo MP, Di Virgilio F, Adinolfi E (2019) The P2X7 receptor modulates immune cells infiltration, ectonucleotidases expression and extracellular ATP levels in the tumor microenvironment. *Oncogene* 38: 3636–3650
- De Sousa EMF, Vermeulen L, Fessler E, Medema JP (2013) Cancer heterogeneity—a multifaceted view. *EMBO Rep* 14: 686–695
- Deng WW, Li YC, Ma SR, Mao L, Yu GT, Bu LL, Kulkarni AB, Zhang WF, Sun ZJ (2018) Specific blockade CD73 alters the "exhausted" phenotype of T cells in head and neck squamous cell carcinoma. *Int J Cancer* 143: 1494–1504
- Dersh D, Hollý J, Yewdell JW (2021) A few good peptides: MHC class I-based cancer immunosurveillance and immunoevasion. *Nat Rev Immunol* 21: 116–128
- Di Virgilio F, Sarti AC, Falzoni S, De Marchi E, Adinolfi E (2018) Extracellular ATP and P2 purinergic signalling in the tumour microenvironment. *Nat Rev Cancer* 18: 601–618
- Dosch M, Gerber J, Jebbawi F, Beldi G (2018) Mechanisms of ATP release by inflammatory cells. *Int J Mol Sci* 19: 1222
- Dosch M, Zindel J, Jebbawi F, Melin N, Sanchez-Taltavull D, Stroka D, Candinas D, Beldi G (2019) Connexin-43-dependent ATP release mediates macrophage activation during sepsis. *Elife* 8: e42670
- Douguet L, Janho dit Hreich S, Benzaquen J, Seguin L, Juhel T, Dezitter X, Duranton C, Ryffel B, Kanellopoulos J, Delarasse C et al (2021) A small-molecule P2RX7 activator promotes anti-tumor immune responses and sensitizes lung tumor to immunotherapy. *Nat Commun* 12: 653
- Duan S, Nordmeier S, Byrnes AE, Buxton ILO (2021) Extracellular vesicle-mediated purinergic signaling contributes to host microenvironment plasticity and metastasis in triple negative breast cancer. *Int J Mol Sci* 22: 597

- Dunn GP, Bruce AT, Ikeda H, Old LJ, Schreiber RD (2002) Cancer immunoeediting: from immunosurveillance to tumor escape. *Nat Immunol* 3: 991–998
- Elliott MR, Chekeni FB, Trampont PC, Lazarowski ER, Kadl A, Walk SF, Park D, Woodson RI, Ostankovich M, Sharma P et al (2009) Nucleotides released by apoptotic cells act as a find-me signal to promote phagocytic clearance. *Nature* 461: 282–286
- Eltzschig HK, Eckle T, Mager A, Küper N, Karcher C, Weissmüller T, Boengler K, Schulz R, Robson SC, Colgan SP (2006) ATP release from activated neutrophils occurs via connexin 43 and modulates adenosine-dependent endothelial cell function. *Circ Res* 99: 1100–1108
- Facon T, Kumar S, Plesner T, Orłowski RZ, Moreau P, Bahlis N, Basu S, Nahi H, Hulin C, Quach H et al (2019) Daratumumab plus Lenalidomide and Dexamethasone for Untreated Myeloma. *N Engl J Med* 380: 2104–2115
- Fong L, Hotson A, Powderly JD, Sznol M, Heist RS, Choueiri TK, George S, Hughes BGM, Hellmann MD, Shepard DR et al (2020) Adenosine 2A receptor blockade as an immunotherapy for treatment-refractory renal cell cancer. *Cancer Discov* 10: 40–53
- Fridman WH, Zitvogel L, Sautès-Fridman C, Kroemer G (2017) The immune contexture in cancer prognosis and treatment. *Nat Rev Clin Oncol* 14: 717–734
- Gaertner F, Massberg S (2019) Patrolling the vascular borders: platelets in immunity to infection and cancer. *Nat Rev Immunol* 19: 747–760
- Gallerano D, Ciminati S, Grimaldi A, Piconese S, Cammarata I, Focaccetti C, Pacella I, Accapezzato D, Lancellotti F, Sacco L et al (2020) Genetically driven CD39 expression shapes human tumor-infiltrating CD8(+) T-cell functions. *Int J Cancer* 147: 2597–2610
- Galluzzi L, Chan TA, Kroemer G, Wolchok JD, Lopez-Soto A (2018) The hallmarks of successful anticancer immunotherapy. *Sci Transl Med* 10: eaat7807
- Galluzzi L, Humeau J, Buqué A, Zitvogel L, Kroemer G (2020) Immunostimulation with chemotherapy in the era of immune checkpoint inhibitors. *Nat Rev Clin Oncol* 17: 725–741
- Gao ZW, Wang HP, Lin F, Wang X, Long M, Zhang HZ, Dong K (2017) CD73 promotes proliferation and migration of human cervical cancer cells independent of its enzyme activity. *BMC Cancer* 17: 135
- Garner H, de Visser KE (2020) Immune crosstalk in cancer progression and metastatic spread: a complex conversation. *Nat Rev Immunol* 20: 483–497
- Ghiringhelli F, Apetoh L, Tesniere A, Aymeric L, Ma Y, Ortiz C, Vermaelen K, Panaretakis T, Mignot G, Ullrich E et al (2009) Activation of the NLRP3 inflammasome in dendritic cells induces IL-1 β -dependent adaptive immunity against tumors. *Nat Med* 15: 1170–1178
- Giatromanolaki A, Kouroupi M, Pouliou S, Mitrakas A, Hasan F, Pappa A, Koukourakis MI (2020) Ectonucleotidase CD73 and CD39 expression in non-small cell lung cancer relates to hypoxia and immunosuppressive pathways. *Life Sci* 259: 118389
- Goswami S, Walle T, Cornish AE, Basu S, Anandhan S, Fernandez I, Vence L, Blando J, Zhao H, Yadav SS et al (2020) Immune profiling of human tumors identifies CD73 as a combinatorial target in glioblastoma. *Nat Med* 26: 39–46
- Hanahan D, Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell* 144: 646–674
- Hatfield SM, Kjaergaard J, Lukashev D, Belikoff B, Schreiber TH, Sethumadhavan S, Abbott R, Philbrook P, Thayer M, Shujia D et al (2014) Systemic oxygenation weakens the hypoxia and hypoxia inducible factor 1 α -dependent and extracellular adenosine-mediated tumor protection. *J Mol Med (Berl)* 92: 1283–1292
- Hatfield SM, Sitkovsky MV (2020) Antihypoxic oxygenation agents with respiratory hyperoxia to improve cancer immunotherapy. *J Clin Invest* 130: 5629–5637
- Häusler SF, Del Barrio IM, Diessner J, Stein RG, Strohschein J, Hönig A, Dietl J, Wischhusen J (2014) Anti-CD39 and anti-CD73 antibodies A1 and 7G2 improve targeted therapy in ovarian cancer by blocking adenosine-dependent immune evasion. *Am J Transl Res* 6: 129–139
- Häusler SFM, Montalbán del Barrio I, Strohschein J, Anoop Chandran P, Engel JB, Hönig A, Ossadnik M, Horn E, Fischer B, Krockenberger M et al (2011) Ectonucleotidases CD39 and CD73 on OvCA cells are potent adenosine-generating enzymes responsible for adenosine receptor 2A-dependent suppression of T cell function and NK cell cytotoxicity. *Cancer Immunol Immunother* 60: 1405–1418
- Hay CM, Sult E, Huang Q, Mulgrew K, Fuhrmann SR, McGlinchey KA, Hammond SA, Rothstein R, Rios-Doria J, Poon E et al (2016) Targeting CD73 in the tumor microenvironment with MEDI9447. *Oncoimmunology* 5: e1208875
- Ho WJ, Jaffee EM, Zheng L (2020) The tumour microenvironment in pancreatic cancer - clinical challenges and opportunities. *Nat Rev Clin Oncol* 17: 527–540
- Husain Z, Huang Y, Seth P, Sukhatme VP (2013) Tumor-derived lactate modifies antitumor immune response: effect on myeloid-derived suppressor cells and NK cells. *J Immunol* 191: 1486–1495
- Iannone R, Miele L, Maiolino P, Pinto A, Morello S (2014) Adenosine limits the therapeutic effectiveness of anti-CTLA4 mAb in a mouse melanoma model. *Am J Cancer Res* 4: 172–181
- Imura Y, Morizawa Y, Komatsu R, Shibata K, Shinozaki Y, Kasai H, Moriishi K, Moriyama Y, Koizumi S (2013) Microglia release ATP by exocytosis. *Glia* 61: 1320–1330
- Jiang L, Huang S, Li W, Zhang D, Zhang S, Zhang W, Zheng P, Chen Z (2012) Expression of autophagy and ER stress-related proteins in primary salivary adenoid cystic carcinoma. *Pathol Res Pract* 208: 635–641
- Kamai T, Kijima T, Tsuzuki T, Nukui A, Abe H, Arai K, Yoshida KI (2021) Increased expression of adenosine 2A receptors in metastatic renal cell carcinoma is associated with poorer response to anti-vascular endothelial growth factor agents and anti-PD-1/Anti-CTLA4 antibodies and shorter survival. *Cancer Immunol Immunother*
- Kamata-Sakurai M, Narita Y, Hori Y, Nemoto T, Uchikawa R, Honda M, Hironiwa N, Taniguchi K, Shida-Kawazoe M, Metsugi S et al (2020) Antibody to CD137 activated by extracellular adenosine triphosphate is tumor selective and broadly effective in vivo without systemic immune activation. *Cancer Discov* 11: 158–175
- Kang J, Kang N, Lovatt D, Torres A, Zhao Z, Lin J, Nedergaard M (2008) Connexin 43 hemichannels are permeable to ATP. *J Neurosci* 28: 4702–4711
- Kashyap AS, Thelemann T, Klar R, Kallert SM, Festag J, Buchi M, Hinterwimmer L, Schell M, Michel S, Jaschinski F et al (2019) Antisense oligonucleotide targeting CD39 improves anti-tumor T cell immunity. *J Immunother Cancer* 7: 67
- Kayagaki N, Kornfeld OS, Lee BL, Stowe IB, O'Rourke K, Li Q, Sandoval W, Yan D, Kang J, Xu M et al (2021) NINJ1 mediates plasma membrane rupture during lytic cell death. *Nature* 591: 131–136
- Kayhan M, Koyas A, Akdemir I, Savas AC, Cekic C (2019) Adenosine receptor signaling targets both PKA and Epac pathways to polarize dendritic cells to a suppressive phenotype. *J Immunol* 203: 3247–3255
- Kepp O, Kroemer G (2020) Autophagy induction by thiostrepton for the improvement of anticancer therapy. *Autophagy* 16: 1166–1167
- Kim JS, Bae GE, Kim KH, Lee SI, Chung C, Lee D, Lee TH, Kwon IS, Yeo MK (2019) Prognostic significance of LC3B and p62/SQSTM1 expression in gastric adenocarcinoma. *Anticancer Res* 39: 6711–6722
- Kjaergaard J, Hatfield S, Jones G, Ohta A, Sitkovsky M (2018) A(2A) adenosine receptor gene deletion or synthetic A(2A) antagonist liberate

- tumor-reactive CD8(+) T cells from tumor-induced immunosuppression. *J Immunol* 201: 782–791
- Klein M, Bopp T (2016) Cyclic AMP represents a crucial component of Treg cell-mediated immune regulation. *Front Immunol* 7: 315
- Ko YH, Cho YS, Won HS, Jeon EK, An HJ, Hong SU, Park JH, Lee MA (2013) Prognostic significance of autophagy-related protein expression in resected pancreatic ductal adenocarcinoma. *Pancreas* 42: 829–835
- Koh J, Kim Y, Lee KY, Hur JY, Kim MS, Kim B, Cho HJ, Lee YC, Bae YH, Ku BM et al (2020) MDSC subtypes and CD39 expression on CD8(+) T cells predict the efficacy of anti-PD-1 immunotherapy in patients with advanced NSCLC. *Eur J Immunol* 50: 1810–1819
- Kroemer G, Senovilla L, Galluzzi L, André F, Zitvogel L (2015) Natural and therapy-induced immunosurveillance in breast cancer. *Nat Med* 21: 1128–1138
- Krstic J, Pieber TR, Prokesch A (2020) Stratifying nutritional restriction in cancer therapy: next stop, personalized medicine. *Int Rev Cell Mol Biol* 354: 231–259
- Künzli BM, Berberat PO, Giese T, Csizmadia E, Kaczmarek E, Baker C, Halaceli I, Büchler MW, Friess H, Robson SC (2007) Upregulation of CD39/NTPDases and P2 receptors in human pancreatic disease. *Am J Physiol Gastrointest Liver Physiol* 292: G223–G230
- Ladoire S, Enot D, Senovilla L, Ghiringhelli F, Poirier-Colame V, Chaba K, Semeraro M, Chaix M, Penault-Llorca F, Arnould L et al (2016) The presence of LC3B puncta and HMGB1 expression in malignant cells correlate with the immune infiltrate in breast cancer. *Autophagy* 12: 864–875
- Ladoire S, Penault-Llorca F, Senovilla L, Dalban C, Enot D, Locher C, Prada N, Poirier-Colame V, Chaba K, Arnould L et al (2015) Combined evaluation of LC3B puncta and HMGB1 expression predicts residual risk of relapse after adjuvant chemotherapy in breast cancer. *Autophagy* 11: 1878–1890
- Lan J, Lu H, Samanta D, Salman S, Lu Y, Semenza GL (2018) Hypoxia-inducible factor 1-dependent expression of adenosine receptor 2B promotes breast cancer stem cell enrichment. *Proc Natl Acad Sci USA* 115: E9640–E9648
- Lapps CM, Rieger JM, Linden J (2005) A2A adenosine receptor induction inhibits IFN- γ production in murine CD4+ T cells. *J Immunol* 174: 1073–1080
- Leclerc BC, Charlebois R, Chouinard G, Allard B, Pommey S, Saad F, Stagg J (2016) CD73 expression is an independent prognostic factor in prostate cancer. *Clin Cancer Res* 22: 158–166
- Lee YJ, Ha YJ, Na Kang Yu, Kang KJ, Hwang JS, Chung WJ, Cho KB, Park KS, Kim ES, Seo H-Y et al (2013) The autophagy-related marker LC3 can predict prognosis in human hepatocellular carcinoma. *PLoS One* 8: e81540
- Lee YS, Radford KJ (2019) The role of dendritic cells in cancer. *Int Rev Cell Mol Biol* 348: 123–178
- Leone RD, Powell JD (2020) Metabolism of immune cells in cancer. *Nat Rev Cancer* 20: 516–531
- Leone RD, Sun IM, Oh MH, Sun IH, Wen J, Englert J, Powell JD (2018) Inhibition of the adenosine A2a receptor modulates expression of T cell coinhibitory receptors and improves effector function for enhanced checkpoint blockade and ACT in murine cancer models. *Cancer Immunol Immunother* 67: 1271–1284
- Li C, Chen Q, Tian Z, Li S, Gong Z, Lin Z, Wang B, Liu H (2019a) Expression of MIF, Beclin1, and LC3 in human salivary gland adenoid cystic carcinoma and its prognostic value. *Medicine* 98: e15402
- Li H, Lv M, Qiao B, Li X (2019b) Blockade of CD73/adenosine axis improves the therapeutic efficacy of docetaxel in epithelial ovarian cancer. *Arch Gynecol Obstet* 299: 1737–1746
- Li J, Wang L, Chen X, Li L, Li Yu, Ping Yu, Huang L, Yue D, Zhang Z, Wang F et al (2017) CD39/CD73 upregulation on myeloid-derived suppressor cells via TGF- β -mTOR-HIF-1 signaling in patients with non-small cell lung cancer. *Oncotarget* 6: e1320011
- Li JX, Yan Q, Liu N, Zheng WJ, Hu M, Yu ZM, Zhou YD, Wang XW, Liang FX, Chen R (2020) The prognostic value of autophagy-related markers Bcln-1 and LC-3 in colorectal cancers: a systematic review and meta-analysis. *Evid Based Complement Alternat Med* 2020: 8475840
- Li X-Y, Moesta AK, Xiao C, Nakamura K, Casey M, Zhang H, Madore J, Lepletier A, Aguilera AR, Sundarrajan A et al (2019c) Targeting CD39 in cancer reveals an extracellular ATP- and inflammasome-driven tumor immunity. *Cancer Discov* 9: 1754–1773
- Lim EA, Bauer TM, Patel MR, Falchook GS, Karlix JL, Choe JH, George DJ, Mugundu GM, Pilling E, Chen H et al (2020) A phase I, open-label, multicenter study to assess the safety, pharmacokinetics, and preliminary antitumor activity of AZD4635 both as monotherapy and in combination in patients with advanced solid malignancies: Results from prostate cancer patients (NCT02740985). *J Clin Oncol* 38: 5518
- Lim To WK, Kumar P, Marshall JM (2015) Hypoxia is an effective stimulus for vesicular release of ATP from human umbilical vein endothelial cells. *Placenta* 36: 759–766
- Loi S, Pommey S, Haibe-Kains B, Beavis PA, Darcy PK, Smyth MJ, Stagg J (2013) CD73 promotes anthracycline resistance and poor prognosis in triple negative breast cancer. *Proc Natl Acad Sci USA* 110: 11091–11096
- Lopez-Otin C, Kroemer G (2021) Hallmarks of health. *Cell* 184: 33–63
- Ludwig N, Yerneni SS, Azambuja JH, Gillespie DG, Menshikova EV, Jackson EK, Whiteside TL (2020) Tumor-derived exosomes promote angiogenesis via adenosine A(2B) receptor signaling. *Angiogenesis* 23: 599–610
- Ma Y, Adjemian S, Mattarollo S, Yamazaki T, Aymeric L, Yang H, Portela Catani J, Hannani D, Duret H, Steegh K et al (2013) Anticancer chemotherapy-induced intratumoral recruitment and differentiation of antigen-presenting cells. *Immunity* 38: 729–741
- Ma Y, Aymeric L, Locher C, Mattarollo SR, Delahaye NF, Pereira P, Boucontet L, Apetoh L, Ghiringhelli F, Casares N et al (2011) Contribution of IL-17-producing gamma delta T cells to the efficacy of anticancer chemotherapy. *J Exp Med* 208: 491–503
- Mantovani A, Marchesi F, Malesci A, Laghi L, Allavena P (2017) Tumour-associated macrophages as treatment targets in oncology. *Nat Rev Clin Oncol* 14: 399–416
- Martin S, Dudek-Peric AM, Garg AD, Roose H, Demirsoy S, Van Eygen S, Mertens F, Vangheluwe P, Vankelecom H, Agostinis P (2017) An autophagy-driven pathway of ATP secretion supports the aggressive phenotype of BRAF(V600E) inhibitor-resistant metastatic melanoma cells. *Autophagy* 13: 1512–1527
- Martinek J, Wu TC, Cadena D, Banchereau J, Palucka K (2019) Interplay between dendritic cells and cancer cells. *Int Rev Cell Mol Biol* 348: 179–215
- Martinez-Outschoorn UE, Peiris-Pagés M, Pestell RG, Sotgia F, Lisanti MP (2017) Cancer metabolism: a therapeutic perspective. *Nat Rev Clin Oncol* 14: 113
- Martins I, Wang Y, Michaud M, Ma Y, Sukkurwala AQ, Shen S, Kepp O, Métiévier D, Galluzzi L, Perfettini J-I et al (2014) Molecular mechanisms of ATP secretion during immunogenic cell death. *Cell Death Differ* 21: 79–91
- Medina CB, Mehrotra P, Arandjelovic S, Perry JSA, Guo Y, Morioka S, Barron B, Walk SF, Ghesquière B, Krupnick AS et al (2020) Metabolites released from apoptotic cells act as tissue messengers. *Nature* 580: 130–135
- Medler T, Patel JM, Alice A, Baird JR, Hu HM, Gough MJ (2019) Activating the nucleic acid-sensing machinery for anticancer immunity. *Int Rev Cell Mol Biol* 344: 173–214

- Messaoudi N, Cousineau I, Arslanian E, Henault D, Stephen D, Vandenbroucke-Menu F, Dagenais M, Létourneau R, Plasse M, Roy A et al (2020) Prognostic value of CD73 expression in resected colorectal cancer liver metastasis. *Oncoimmunology* 9: 1746138
- Michaud M, Martins I, Sukkurwala Aq, Adjemian S, Ma Y, Pellegatti P, Shen S, Kepp O, Scoazec M, Mignot G et al (2011) Autophagy-dependent anticancer immune responses induced by chemotherapeutic agents in mice. *Science* 334: 1573–1577
- Minguet S, Huber M, Rosenkranz L, Schamel WW, Reth M, Brummer T (2005) Adenosine and cAMP are potent inhibitors of the NF- κ B pathway downstream of immunoreceptors. *Eur J Immunol* 35: 31–41
- Mittal D, Sinha D, Barkauskas D, Young A, Kalimutho M, Stannard K, Caramia F, Haibe-Kains B, Stagg J, Khanna KK et al (2016) Adenosine 2B receptor expression on cancer cells promotes metastasis. *Cancer Res* 76: 4372–4382
- Mittal D, Young A, Stannard K, Yong M, Teng MW, Allard B, Stagg J, Smyth MJ (2014) Antimetastatic effects of blocking PD-1 and the adenosine A2A receptor. *Cancer Res* 74: 3652–3658
- Mobasher M, Miller RA, Kwei L, Strahs D, Das V, Luciano G, Powderly JD, Merchan JR, Barve MA, LoRusso P et al (2019) A phase I/II multicenter study to evaluate the humanized anti-CD73 antibody, CPI-006, as a single agent, in combination with CPI-444, and in combination with pembrolizumab in adult patients with advanced cancers. *J Clin Oncol* 37: TPS2646-TPS2646
- Moesta AK, Li XY, Smyth MJ (2020) Targeting CD39 in cancer. *Nat Rev Immunol* 20: 739–755
- Morandi F, Morandi B, Horenstein AL, Chillemi A, Quarona V, Zaccarello G, Carrega P, Ferlazzo G, Mingari MC, Moretta L et al (2015) A non-canonical adenosinergic pathway led by CD38 in human melanoma cells induces suppression of T cell proliferation. *Oncotarget* 6: 25602–25618
- Murata N, Ito S, Furuya K, Takahara N, Naruse K, Aso H, Kondo M, Sokabe M, Hasegawa Y (2014) Ca²⁺ influx and ATP release mediated by mechanical stretch in human lung fibroblasts. *Biochem Biophys Res Commun* 453: 101–105
- Nagata Y, Ezo S, Fujita J, Okuzaki D, Motooka D, Ishibashi T, Ichii M, Tanimura A, Kurashige M, Morii E et al (2021) Ectonucleotidase CD39 is highly expressed on ATLL cells and is responsible for their immunosuppressive function. *Leukemia* 35: 107–118
- Nakamura K, Casey M, Oey H, Vari F, Stagg J, Gandhi MK, Smyth MJ (2020) Targeting an adenosine-mediated "don't eat me signal" augments anti-lymphoma immunity by anti-CD20 monoclonal antibody. *Leukemia* 34: 2708–2721
- Narahari AK, Kreutzberger AJ, Gaete PS, Chiu YH, Leonhardt SA, Medina CB, Jin X, Oleniacz PW, Kiessling V, Barrett PQ et al (2021) ATP and large signaling metabolites flux through caspase-activated Pannexin 1 channels. *Elife* 10: e64787
- Németh ZH, Lutz CS, Csóka B, Deitch EA, Leibovich SJ, Gause WC, Tone M, Pacher P, Vizi ES, Haskó G (2005) Adenosine augments IL-10 production by macrophages through an A2B receptor-mediated posttranscriptional mechanism. *J Immunol* 175: 8260–8270
- Neo SY, Yang Y, Record J, Ma R, Chen X, Chen Z, Tobin NP, Blake E, Seitz C, Thomas R et al (2020) CD73 immune checkpoint defines regulatory NK cells within the tumor microenvironment. *J Clin Invest* 130: 1185–1198
- Nüchel H, Frey UH, Dürig J, Dührsen U, Siffert W (2004) 1513A/C polymorphism in the P2X7 receptor gene in chronic lymphocytic leukemia: absence of correlation with clinical outcome. *Eur J Haematol* 72: 259–263
- Ohta A, Gorelik E, Prasad SJ, Ronchese F, Lukaszew D, Wong M, Huang X, Caldwell S, Liu K, Smith P et al (2006) A2A adenosine receptor protects tumors from antitumor T cells. *Proc Natl Acad Sci USA* 103: 13132–13137
- O'Sullivan D, Sanin DE, Pearce EJ, Pearce EL (2019) Metabolic interventions in the immune response to cancer. *Nat Rev Immunol* 19: 324–335
- Paneesha S, Starczynski J, Pepper C, Delgado J, Hooper L, Fegan C, Pratt G (2006) The P2X7 receptor gene polymorphism 1513 A→C has no effect on clinical prognostic markers and survival in multiple myeloma. *Leuk Lymphoma* 47: 281–284
- Pellegatti P, Raffaghello L, Bianchi G, Piccardi F, Pistoia V, Di Virgilio F (2008) Increased level of extracellular ATP at tumor sites: in vivo imaging with plasma membrane luciferase. *PLoS One* 3: e2599
- Perregaux DG, McNiff P, Laliberte R, Conklyn M, Gabel CA (2000) ATP acts as an agonist to promote stimulus-induced secretion of IL-1 beta and IL-18 in human blood. *J Immunol* 165: 4615–4623
- Perrot I, Michaud H-A, Giraudon-Paoli M, Augier S, Docquier A, Gros L, Courtois R, Déjou C, Jecko D, Becquart O et al (2019) Blocking antibodies targeting the CD39/CD73 immunosuppressive pathway unleash immune responses in combination cancer therapies. *Cell Rep* 27: 2411–2425
- Perry C, Hazan-Halevy I, Kay S, Cipok M, Grisar D, Deutsch V, Polliack A, Naparstek E, Herishanu Y (2012) Increased CD39 expression on CD4(+) T lymphocytes has clinical and prognostic significance in chronic lymphocytic leukemia. *Ann Hematol* 91: 1271–1279
- Petroni G, Buque A, Zitvogel L, Kroemer G, Galluzzi L (2021) Immunomodulation by targeted anticancer agents. *Cancer Cell* 39: 310–345
- Pietrocola F, Pol J, Vacchelli E, Rao S, Enot D, Baracco E, Levesque S, Castoldi F, Jacquelot N, Yamazaki T et al (2016) Caloric restriction mimetics enhance anticancer immunosurveillance. *Cancer Cell* 30: 147–160
- Pinheiro AR, Paramos-de-Carvalho D, Certal M, Costa MA, Costa C, Magalhães-Cardoso MT, Ferreirinha F, Sévigny J, Correia-de-Sá P (2013) Histamine induces ATP release from human subcutaneous fibroblasts, via pannexin-1 hemichannels, leading to Ca²⁺ mobilization and cell proliferation. *J Biol Chem* 288: 27571–27583
- Pulte D, Furman RR, Broekman MJ, Drosopoulos JH, Ballard HS, Olson KE, Kizer JR, Marcus AJ (2011) CD39 expression on T lymphocytes correlates with severity of disease in patients with chronic lymphocytic leukemia. *Clin Lymphoma Myeloma Leuk* 11: 367–372
- Qi Yu, Xia Yu, Lin Z, Qu Y, Qi Y, Chen Y, Zhou Q, Zeng H, Wang J, Chang Y et al (2020) Tumor-infiltrating CD39(+)CD8(+) T cells determine poor prognosis and immune evasion in clear cell renal cell carcinoma patients. *Cancer Immunol Immunother* 69: 1565–1576
- Qin Z, Yu X, Lin M, Wu J, Ma S, Wang N (2018) Prognostic and clinicopathological value of Beclin-1 expression in hepatocellular carcinoma: a meta-analysis. *World J Surg Oncol* 16: 170
- Rao S, Gharib K, Han A (2019) Cancer immunosurveillance by T cells. *Int Rev Cell Mol Biol* 342: 149–173
- Raskovalova T, Huang X, Sitkovsky M, Zacharia LC, Jackson EK, Gorelik E (2005) Gs protein-coupled adenosine receptor signaling and lytic function of activated NK cells. *J Immunol* 175: 4383–4391
- Rodriguez-Ruiz ME, Vitale I, Harrington KJ, Melero I, Galluzzi L (2020) Immunological impact of cell death signaling driven by radiation on the tumor microenvironment. *Nat Immunol* 21: 120–134
- Rybstein MD, Bravo-San Pedro JM, Kroemer G, Galluzzi L (2018) The autophagic network and cancer. *Nat Cell Biol* 20: 243–251
- Sáez JC, Contreras-Duarte S, Gómez GI, Labra VC, Santibañez CA, Gajardo-Gómez R, Avendaño BC, Díaz EF, Montero TD, Velarde V et al (2018) Connexin 43 hemichannel activity promoted by pro-inflammatory cytokines and high glucose alters endothelial cell function. *Front Immunol* 9: 1899

- Saez PJ, Vargas P, Shoji KF, Harcha PA, Lennon-Dumenil AM, Saez JC (2017) ATP promotes the fast migration of dendritic cells through the activity of pannexin 1 channels and P2X7 receptors. *Sci Signal* 10: eaah7107
- Saleh A, Simone B, Palazzo J, Savage JE, Sano Y, Dan Tu, Jin L, Champ C, Zhao S, Lim M et al (2013) Caloric restriction augments radiation efficacy in breast cancer. *Cell Cycle* 12: 1955–1963
- Schuler PJ, Saze Z, Hong CS, Muller L, Gillespie DG, Cheng D, Harasymczuk M, Mandapathil M, Lang S, Jackson EK et al (2014) Human CD4⁺ CD39⁺ regulatory T cells produce adenosine upon co-expression of surface CD73 or contact with CD73⁺ exosomes or CD73⁺ cells. *Clin Exp Immunol* 177: 531–543
- Schumacher D, Strlic B, Sivaraj KK, Wettschureck N, Offermanns S (2013) Platelet-derived nucleotides promote tumor-cell transendothelial migration and metastasis via P2Y2 receptor. *Cancer Cell* 24: 130–137
- Shi L, Feng M, Du S, Wei X, Song H, Yixin X, Song J, Wenxian G (2019) Adenosine generated by regulatory T cells induces CD8(+) T cell exhaustion in gastric cancer through A2aR pathway. *Biomed Res Int* 2019: 4093214
- Sidders B, Zhang P, Goodwin K, O'Connor G, Russell DL, Borodovsky A, Armenia J, McEwen R, Linghu B, Bendell JC et al (2020) Adenosine signaling is prognostic for cancer outcome and has predictive utility for immunotherapeutic response. *Clin Cancer Res* 26: 2176–2187
- Simone BA, Dan Tu, Palagani A, Jin L, Han SY, Wright C, Savage JE, Gitman R, Lim MK, Palazzo J et al (2016) Caloric restriction coupled with radiation decreases metastatic burden in triple negative breast cancer. *Cell Cycle* 15: 2265–2274
- Sitkovsky MV, Hatfield S, Abbott R, Belikoff B, Lukashev D, Ohta A (2014) Hostile, hypoxia-A2-adenosinergic tumor biology as the next barrier to overcome for tumor immunologists. *Cancer Immunol Res* 2: 598–605
- Smyth LA, Ratnasothy K, Tsang JY, Boardman D, Warley A, Lechler R, Lombardi G (2013) CD73 expression on extracellular vesicles derived from CD4⁺ CD25⁺ Foxp3⁺ T cells contributes to their regulatory function. *Eur J Immunol* 43: 2430–2440
- Sprooten J, Agostinis P, Garg AD (2019) Type I interferons and dendritic cells in cancer immunotherapy. *Int Rev Cell Mol Biol* 348: 217–262
- Stagg J, Divisekera U, McLaughlin N, Sharkey J, Pommey S, Denoyer D, Dwyer KM, Smyth MJ (2010) Anti-CD73 antibody therapy inhibits breast tumor growth and metastasis. *Proc Natl Acad Sci USA* 107: 1547–1552
- Stagg J, Divisekera U, Duret H, Sparwasser T, Teng MW, Darcy PK, Smyth MJ (2011) CD73-deficient mice have increased antitumor immunity and are resistant to experimental metastasis. *Cancer Res* 71: 2892–2900
- Stagg J, Smyth MJ (2010) Extracellular adenosine triphosphate and adenosine in cancer. *Oncogene* 29: 5346–5358
- Starzynski J, Pepper C, Pratt G, Hooper L, Thomas A, Hoy T, Milligan D, Bentley P, Fegan C (2003) The P2X7 receptor gene polymorphism 1513 A→C has no effect on clinical prognostic markers, in vitro sensitivity to fludarabine, Bcl-2 family protein expression or survival in B-cell chronic lymphocytic leukaemia. *Br J Haematol* 123: 66–71
- Stout CE, Costantin JL, Naus CC, Charles AC (2002) Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels. *J Biol Chem* 277: 10482–10488
- Sun X, Wu Y, Gao W, Enjyoji K, Cszimadia E, Müller CE, Murakami T, Robson SC (2010) CD39/ENTPD1 expression by CD4⁺Foxp3⁺ regulatory T cells promotes hepatic metastatic tumor growth in mice. *Gastroenterology* 139: 1030–1040
- Synnestvedt K, Furuta GT, Comerford KM, Louis N, Karhausen J, Eltzschig HK, Hansen KR, Thompson LF, Colgan SP (2002) Ecto-5'-nucleotidase (CD73) regulation by hypoxia-inducible factor-1 mediates permeability changes in intestinal epithelia. *J Clin Invest* 110: 993–1002
- Tahkola K, Ahtiainen M, Kellokumpu I, Mecklin JP, Laukkanen J, Laakkonen J, Kenessey I, Jalkanen S, Salmi M, Bohm J (2021) Prognostic impact of CD73 expression and its relationship to PD-L1 in patients with radically treated pancreatic cancer. *Virchows Arch* 478: 209–217
- Talmadge JE, Gabrilovich DI (2013) History of myeloid-derived suppressor cells. *Nat Rev Cancer* 13: 739–752
- Tang H, Sebti S, Titone R, Zhou Y, Isidoro C, Ross TS, Hibshoosh H, Xiao G, Packer M, Xie Y et al (2015) Decreased BECN1 mRNA expression in human breast cancer is associated with estrogen receptor-negative subtypes and poor prognosis. *EBioMedicine* 2: 255–263
- Tappe KA, Budida R, Stankov MV, Frenz T, R Shah H, Volz A, Sutter G, Kalinke U, Behrens GMN (2018) Immunogenic cell death of dendritic cells following modified vaccinia virus Ankara infection enhances CD8(+) T cell proliferation. *Eur J Immunol* 48: 2042–2054
- Tatsuno K, Yamazaki T, Hanlon D, Han P, Robinson E, Sobolev O, Yurter A, Rivera-Molina F, Arshad N, Edelson RL et al (2019) Extracorporeal photochemotherapy induces bona fide immunogenic cell death. *Cell Death Dis* 10: 578
- Thunberg U, Tobin G, Johnson A, Söderberg O, Padyukov L, Hultdin M, Klareskog L, Enblad G, Sundström C, Roos G et al (2002) Polymorphism in the P2X7 receptor gene and survival in chronic lymphocytic leukaemia. *Lancet* 360: 1935–1939
- Timp W, Feinberg AP (2013) Cancer as a dysregulated epigenome allowing cellular growth advantage at the expense of the host. *Nat Rev Cancer* 13: 497–510
- Togashi Y, Shitara K, Nishikawa H (2019) Regulatory T cells in cancer immunosuppression - implications for anticancer therapy. *Nat Rev Clin Oncol* 16: 356–371
- Tokunaga A, Tsukimoto M, Harada H, Moriyama Y, Kojima S (2010) Involvement of SLC17A9-dependent vesicular exocytosis in the mechanism of ATP release during T cell activation. *J Biol Chem* 285: 17406–17416
- Tokunaga R, Cao S, Naseem M, Lo JH, Battaglin F, Puccini A, Berger MD, Soni S, Millstein J, Zhang W et al (2019) Prognostic Effect of adenosine-related genetic variants in metastatic colorectal cancer treated with bevacizumab-based chemotherapy. *Clin Colorectal Cancer* 18: e8–e19
- Torres-Pineda DB, Mora-García MdL, García-Rocha R, Hernández-Montes J, Weiss-Steider B, Montesinos-Montesinos JJ, Don-López CA, Marín-Aquino LA, Muñoz-Godínez R, Ibarra LRÁ et al (2020) Adenosine augments the production of IL-10 in cervical cancer cells through interaction with the A (2B) adenosine receptor, resulting in protection against the activity of cytotoxic T cells. *Cytokine* 130: 155082
- Tripathi A, Lin E, Xie W, Flaifel A, Steinharter JA, Stern Gatof EN, Bouchard G, Fleischer JH, Martinez-Chanza N, Gray C et al (2020) Prognostic significance and immune correlates of CD73 expression in renal cell carcinoma. *J Immunother Cancer* 8: e001467
- Tsukui H, Horie H, Koinuma K, Ohzawa H, Sakuma Y, Hosoya Y, Yamaguchi H, Yoshimura K, Lefor AK, Sata N et al (2020) CD73 blockade enhances the local and abscopal effects of radiotherapy in a murine rectal cancer model. *BMC Cancer* 20: 411
- Turcotte M, Spring K, Pommey S, Chouinard G, Cousineau I, George J, Chen GM, Gendoo DMA, Haibe-Kains B, Karn T et al (2015) CD73 is associated with poor prognosis in high-grade serous ovarian cancer. *Cancer Res* 75: 4494–4503
- Turiello R, Capone M, Giannarelli D, Morretta E, Monti MC, Madonna G, Mallardo D, Festino L, Azzaro R, Levesque MP et al (2020) Serum CD73 is a prognostic factor in patients with metastatic melanoma and is associated with response to anti-PD-1 therapy. *J Immunother Cancer* 8: e001689

- Vesely MD, Schreiber RD (2013) Cancer immunoediting: antigens, mechanisms, and implications to cancer immunotherapy. *Ann N Y Acad Sci* 1284: 1–5
- Vitale I, Shema E, Loi S, Galluzzi L (2021) Intratumoral heterogeneity in cancer progression and response to immunotherapy. *Nat Med* 27: 212–224
- Waickman AT, Alme A, Senaldi L, Zarek PE, Horton M, Powell JD (2012) Enhancement of tumor immunotherapy by deletion of the A2A adenosine receptor. *Cancer Immunol Immunother* 61: 917–926
- Wang L, Fan J, Thompson LF, Zhang Y, Shin T, Curriel TJ, Zhang B (2011) CD73 has distinct roles in nonhematopoietic and hematopoietic cells to promote tumor growth in mice. *J Clin Invest* 121: 2371–2382
- Wang M, Jing J, Li H, Liu J, Yuan Y, Sun L (2021) The expression characteristics and prognostic roles of autophagy-related genes in gastric cancer. *PeerJ* 9: e10814
- Wang X, Zhang T, Song Z, Li L, Zhang X, Liu J, Liu X, Qiu L, Qian Z, Zhou S et al (2019) Tumor CD73/A2aR adenosine immunosuppressive axis and tumor-infiltrating lymphocytes in diffuse large B-cell lymphoma: correlations with clinicopathological characteristics and clinical outcome. *Int J Cancer* 145: 1414–1422
- Wang Y, Xie W, Humeau J, Chen G, Liu P, Pol J, Zhang Z, Kepp O, Kroemer G (2020) Autophagy induction by thiostrepton improves the efficacy of immunogenic chemotherapy. *J Immunother Cancer* 8: e000462
- Watson MJ, Vignali PDA, Mullett SJ, Overacre-Delgoffe AE, Peralta RM, Grebinoski S, Menk AV, Rittenhouse NL, DePeaux K, Whetstone RD et al (2021) Metabolic support of tumour-infiltrating regulatory T cells by lactic acid. *Nature* 591: 645–651
- Wennerberg E, Spada S, Rudqvist N-P, Lhuillier C, Gruber S, Chen Q, Zhang F, Zhou XK, Gross SS, Formenti SC et al (2020) CD73 blockade promotes dendritic cell infiltration of irradiated tumors and tumor rejection. *Cancer Immunol Res* 8: 465–478
- Wiley JS, Dao-Ung LP, Gu BJ, Sluyter R, Shemon AN, Li C, Taper J, Gallo J, Manoharan A (2002) A loss-of-function polymorphic mutation in the cytosolic P2X7 receptor gene and chronic lymphocytic leukaemia: a molecular study. *Lancet* 359: 1114–1119
- Willingham SB, Ho PY, Hotson A, Hill C, Piccione EC, Hsieh J, Liu L, Buggy JJ, McCaffery I, Miller RA (2018) A2AR antagonism with CPI-444 induces antitumor responses and augments efficacy to anti-PD-(L)1 and anti-CTLA-4 in preclinical models. *Cancer Immunol Res* 6: 1136–1149
- Wu J, Wang YC, Xu WH, Luo WJ, Wan FN, Zhang HL, Ye DW, Qu YY, Zhu YP (2020a) High expression of CD39 is associated with poor prognosis and immune infiltrates in clear cell renal cell carcinoma. *Onco Targets Ther* 13: 10453–10464
- Wu Qi, Tian A-L, Durand S, Aprahamian F, Nirmalathasan N, Xie W, Liu P, Zhao L, Zhang S, Pan H et al (2020b) Isobacacalchone induces autophagy and improves the outcome of immunogenic chemotherapy. *Cell Death Dis* 11: 1015
- Xiao Z, Dai Z, Locasale JW (2019) Metabolic landscape of the tumor microenvironment at single cell resolution. *Nat Commun* 10: 3763
- Xu S, Shao Q-Q, Sun J-T, Yang N, Xie Qi, Wang D-H, Huang Q-B, Huang B, Wang X-Y, Li X-G et al (2013) Synergy between the ectoenzymes CD39 and CD73 contributes to adenosinergic immunosuppression in human malignant gliomas. *Neuro Oncol* 15: 1160–1172
- Yamazaki T, Kirchmair A, Sato Ai, Buqué A, Rybstein M, Petroni G, Bloy N, Finotello F, Stafford L, Navarro Manzano E et al (2020) Mitochondrial DNA drives abscopal responses to radiation that are inhibited by autophagy. *Nat Immunol* 21: 1160–1171
- Yan A, Joachims ML, Thompson LF, Miller AD, Canoll PD, Bynoe MS (2019) CD73 promotes glioblastoma pathogenesis and enhances its chemoresistance via A(2B) adenosine receptor signaling. *J Neurosci* 39: 4387–4402
- Yan J, Li X-Y, Roman Aguilera A, Xiao C, Jacobberger-Foissac C, Nowlan B, Robson SC, Beers C, Moesta AK, Geetha N et al (2020) Control of metastases via myeloid CD39 and NK cell effector function. *Cancer Immunol Res* 8: 356–367
- Yang H, Geng YH, Wang P, Yang H, Zhou YT, Zhang HQ, He HY, Fang WG, Tian XX (2020a) Extracellular ATP promotes breast cancer invasion and chemoresistance via SOX9 signaling. *Oncogene* 39: 5795–5810
- Yang R, Elsaadi S, Misund K, Abdollahi P, Vandsemb EN, Moen SH, Kusnierczyk A, Slupphaug G, Standal T, Waage A et al (2020b) Conversion of ATP to adenosine by CD39 and CD73 in multiple myeloma can be successfully targeted together with adenosine receptor A2A blockade. *J Immunother Cancer* 8: e000610
- Yang X, Lu Y, Hang J, Zhang J, Zhang T, Huo Y, Liu J, Lai S, Luo D, Wang L et al (2020c) Lactate-modulated immunosuppression of myeloid-derived suppressor cells contributes to the radioresistance of pancreatic cancer. *Cancer Immunol Res* 8: 1440–1451
- Yang Y, Delalio LJ, Best AK, Macal E, Milstein J, Donnelly I, Miller AM, McBride M, Shu X, Koval M et al (2020d) Endothelial Pannexin 1 channels control inflammation by regulating intracellular calcium. *J Immunol* 204: 2995–3007
- Yeaman MR (2014) Platelets: at the nexus of antimicrobial defence. *Nat Rev Microbiol* 12: 426–437
- Yegutkin GG, Marttila-Ichihara F, Karikoski M, Niemelä J, Laurila JP, Elima K, Jalkanen S, Salmi M (2011) Altered purinergic signaling in CD73-deficient mice inhibits tumor progression. *Eur J Immunol* 41: 1231–1241
- Young A, Ngiow S, Barkauskas D, Sult E, Hay C, Blake S, Huang Q, Liu J, Takeda K, Teng M et al (2016) Co-inhibition of CD73 and A2AR adenosine signaling improves anti-tumor immune responses. *Cancer Cell* 30: 391–403
- Young A, Ngiow SF, Gao Y, Patch A-M, Barkauskas DS, Messaoudene M, Lin G, Coudert JD, Stannard KA, Zitvogel L et al (2018) A2AR adenosine signaling suppresses natural killer cell maturation in the tumor microenvironment. *Cancer Res* 78: 1003–1016
- Yu M, Guo G, Huang L, Deng L, Chang C-S, Achyut BR, Canning M, Xu N, Arbab AS, Bollag RJ et al (2020) CD73 on cancer-associated fibroblasts enhanced by the A2B-mediated feedforward circuit enforces an immune checkpoint. *Nat Commun* 11: 515
- Zhang LY, Ibbotson RE, Orchard JA, Gardiner AC, Seear RV, Chase AJ, Oscier DG, Cross NC (2003) P2X7 polymorphism and chronic lymphocytic leukaemia: lack of correlation with incidence, survival and abnormalities of chromosome 12. *Leukemia* 17: 2097–2100
- Zhou P, Zhi X, Zhou T, Chen S, Li X, Wang L, Yin L, Shao Z, Ou Z (2007) Overexpression of Ecto-5'-nucleotidase (CD73) promotes T-47D human breast cancer cells invasion and adhesion to extracellular matrix. *Cancer Biol Ther* 6: 426–431